

SHORT ORAL CONTRIBUTIONS (ABSTRACTS)

THE HERBIG-HARO OBJECTS ASSOCIATED WITH ESO 210-6A

J.A. Graham and J.E. Elias

Cerro Tololo Inter-American Observatory, Chile

The observational data are given for the dark globule ESO 210-6A and the associated Herbig-Haro objects HH 46 and HH 47 originally discovered by Schwartz. Images are presented in the light of the H α and [S II] lines. As well, we show the continuum images in the visual, red and near infrared and low resolution infrared contour maps at 1.2, 1.6 and 2.2 microns. We believe that we have located the stellar or pre-stellar source which generates the wind responsible for the HH objects in ESO 210-6A. Velocity data based on spectrograms taken with the CTIO 4m telescope are shown. We discuss the structure and geometry of ESO 210-6A and compare our results with recent observations by Dopita, Evans, and Schwartz.

DISCUSSION

S. Strom: What spectral feature near the "source" led you to deduce a line width of 300 km s^{-1} ?

Graham: The [S II] and [N II] lines were best for this. H α is clearly broad and contaminated by strong radiation from the Gum Nebula.

Tapia: What is the sensitivity limit of your $2.2\mu\text{m}$ survey?

Graham: The lowest contour corresponds to a $2.2\mu\text{m}$ surface brightness of about $19^{\text{th}}5 \text{ (arc sec)}^{-2}$.

IR MAPPING OF HH-OBJECTS 7-11 IN MOLECULAR HYDROGEN EMISSION LINES

W.J. Zealey and P.M. Williams

U.K. Infrared Telescope of the Royal Observatory, Edinburgh, United Kingdom

and

G. Sandell

Observatory and Astrophysical Laboratory
University of Helsinki, Finland

The striking alignment of HH-objects, HH 7-11 is the first group of HH objects to be mapped in the $N = 1-0$, S(1)m $2.12 \mu\text{m}$ line of molecular hydrogen. This mapping was carried out with the 3.8 meter UKIRT at Mauna Kea, Hawaii.

The HH-objects, 7-11 form a chain, aligned with a H $_2$ -IR source. This object appears to be the centre of a high velocity CO outflow and a possible exciting source for the HH-objects.

The molecular hydrogen emission closely follows the extent of the HH chain. It peaks near HH 8 and HH 9. We see a sharp cut off to the east of the HH 7 and to the south of the chain. The emission declines toward the H $_2$ -IR source but can still be detected 10 seconds northwest of it.

A spectrum shows the S(1) and Q branches of the emission to be of similar intensity.

We also detected H $_2$ emission in HH 32A which is associated with the ex-

trème T Tauri star AS353A.

Such high resolution mapping of molecular hydrogen permits a detailed comparison with the optical emission. This represents an important step in understanding the interactions of mass outflows, molecular clouds and HH-objects.

DISCUSSION

Cohen: Are you planning to observe the vicinity of L1551-IRS5 in detail in molecular hydrogen?

Zealey: Our plans for 1983 are to observe the L1551 region. The region is large, therefore we will confine the observations, initially to near the HH-objects. We are also going to observe southern HH/bipolar complexes using the AAT this spring. These sources will include the intriguing HH 46/47 complex.

Snell: Did you map the molecular hydrogen emission in the AS353A region?

Zealey: Yes. We detected the H₂ emission in the 1-0 S(1) 2.12 μ m line, coincident with HH 32A. The surface brightness is 2.6 $\times 10^{-5}$ erg cm⁻² s⁻¹ sr⁻¹. We only observed this single position in the AS353A region.

THE ULTRAVIOLET SPECTRUM OF HERBIG-HARO OBJECT 2H

E.W. Brugel¹, J.M. Shull¹, and C.G. Seab

Laboratory for Atmospheric and Space Physics
University of Colorado, Boulder, CO, USA

We present *IUE* spectra of Herbig-Haro object 2H which show a strong "excess" UV continuum and prominent emission lines of C, N, O, Si, Mg, and possibly Al. The continuum, F λ , exhibits a turnover shortward of about 1450 Å, confirming for the first time the H⁺ 2-photon nature for the emission source. We also note a possible absorption feature near 1680 Å, which could result from a new grain or molecular constituent in these protostellar objects.

Our recently computed models of steady shocks into partially-ionized gas reproduce the 2-photon spectral shape, but its intensity relative to H β and the Balmer continuum is anomalously high. We suggest that a range of shock velocities, 70 to 100 km s⁻¹, or non-steady, "truncated" shocks may be responsible.

1. Guest Investigator with NASA's International Ultraviolet Explorer (*IUE*) telescope.

DISCUSSION

Kuhi: All your conclusions depend on fitting the theoretical two-photon emission to the observed peak in the ultraviolet which looks to me like the blend of several emission lines. How can you rule out this possibility?

Brugel: This is definitely a problem and we cannot completely rule out the possible effects of weak unresolved emission lines. However, the shape of the observed continuum is distinctly characteristic of 2-photon emission!

Böhm: If the extinction is really as small as you indicated, would you not get line ratios in the optical range which are hard to understand using shock wave models?

Brugel: I believe that the average galactic extinction law with E(B-V) = 0.34 is applicable in the optical and the reddening correction is being applied correctly in this wavelength region. However, in the UV the extinction curve is probably much smaller than even the θ^2 Orionis curve. Thus only the UV line strengths need to be adjusted.

Liseau: Three distinct discontinuities were discernible in the observed continuous spectrum of HH 2 ($\lambda \sim 1500, 1700, 2500$ Å). Have you examined the possible continuum contributions from Si I and Mg I ($\lambda \sim 1525, 1680, 2517$ Å)?

Brugel: Only the discontinuity at ~ 1680 Å appears on all 4 images and is probably real. We did consider Mg I continuum and found it to be too weak; we did not consider Si I.

Münch: If the extinction in the object were internal taking into account effects of scattering with high albedo in the visual, it can be shown that the external extinction of the object is nearly zero.

Brugel: Our results also imply very little external extinction.