

## TURBULENT MOTIONS IN MOLECULAR CLOUDS

G.A. Pellegatti Franco, R.D. Tarsia and  
R.J. Quiroga

Observatório Astronômico da Serra da Piedade  
Brasil

ABSTRACT: We have studied the behavior of the inner motions of OH, H<sub>2</sub>CO and CO molecular clouds. This study shows the existence of two main components of these clouds: the narrow one, associated to dense small clouds and a wide one "representing" the large diffuse clouds seen in neutral hydrogen. The large clouds are the "vortex" and intermediate state between turbulent and hydrodynamic motions in the galaxy.

For the dense clouds with sizes  $d < 10\text{pc}$  we have found a relationship  $\sigma \propto d^{0.38}$  consistent with the Kolmogorov law of turbulences; the densities and sizes of these clouds behave as  $n \propto d^{-1}$ . This last relation for these molecular clouds is compared with the H II one. Also, we discuss the effects of the inner magnetic field in these clouds.

*Key words:* Interstellar medium, molecular clouds.

This paper is in press in *Astrophysics and Space Science* (1985).

G.A. Pellegatti Franco, R.J. Quiroga, and R.D. Tarsia: Observatório Astronômico da Serra da Piedade, Departamento de Física, ICEx, UFMG, C.P. 702, 30.000 Belo Horizonte MG Brasil.