

A PRELIMINARY REPORT ON PHOTOMETRIC MEASUREMENTS OF THE
HELIUM LINE $\lambda 10830 \text{ \AA}$

Eugenio E. Mendoza V.
Instituto de Astronomía
Universidad Nacional Autónoma de México

RESUMEN. Este trabajo está basado en ochenta y nueve estrellas, seleccionadas del Catalogo 'Bright Star Catalog'. Las observaciones se han hecho para medir un índice $\psi(25)$, en la misma forma como se definieron los índices $\alpha(16)$ y $\Lambda(9)$. En esta ocasión para medir la absorción total de la línea de helio neutro $\lambda 10830 \text{ \AA}$; se comparan nuestros resultados con las medidas de ancho equivalente determinadas por Zirín. También se efectúan algunas comparaciones con el sistema fotométrico $\alpha(16)\Lambda(9)$.

ABSTRACT. This study is based upon 89 stars taken from the Bright Star Catalog to measure the total absorption of the He I line at $\lambda 10830 \text{ \AA}$. It is defined a $\psi(25)$ index in the same fashion as the $\alpha(16)$ and $\Lambda(9)$ indices. Our results are compared with Zirín's equivalent widths, and with the $\alpha(16)\Lambda(9)$ photometric system.

Key words: PHOTOMETRY

I. INTRODUCTION

Estimates of the He I line at $\lambda 10830 \text{ \AA}$ offers a convenient way of surveying the coronal emission measure of a variety of stars. This has the advantage of being easier and more economical than direct X-ray detection. An extense account of measurements of equivalent widths of this neutral helium line, and applications to Astronomy has been made by Zirín (1982).

This paper gives a description of a photometric technique to measure the He I ($\lambda 10830 \text{ \AA}$) line, in the same fashion as the $\alpha(16)$ and $\Lambda(9)$ indices (Mendoza 1973, 1975, 1979a, 1979b, 1981, 1983, and 1986). Preliminary results are shown.

II. THE OBSERVATIONS

The observations have been carried out with the 1.5 m (Johnson Telescope), and 0.84 m telescopes of the Observatorio Astronomico Nacional at San Pedro Martir, B.C., Mexico in October 1984 and June 1986.

The filters used to define the He I ($\lambda 10830 \text{ \AA}$) index, $\psi(25)$, are described in Table 1, and illustrated graphically in Figure 1. This index is defined as follows:

$$\psi(25) = k + m(b) - 0.5 [m(a) + m(c)] \quad (1)$$

where $m(a)$, $m(b)$, and $m(c)$ are magnitudes measured through the interference filters a,b,c described in Table 1 (see also Fig. 1); k is a constant usually taken as zero. However, herein it will be determined from equivalent widths with the hope that a zero index would mean a negligible equivalent width (see below).


This $\psi(25)$ index is insensitive to the influence of interstellar reddening.

The $\lambda 10830 \text{ \AA}$ line is located in a most difficult spectral region where usually detectors are not very sensitive. We provisionally have employed the same RCA 7102 photomultiplier, used by Johnson to define the RI, and 13-color photometric systems (see per example Johnson and Mitchell 1975). We have observed 89 stars so far, from O to M, spectral types; from I to V, luminosity class; and normal and abnormal stars.

TABLE 1

 $\psi(25)$: He I ($\lambda 10830 \text{ \AA}$) Photometric System

PWL (\AA)	HPWB (\AA)	Transmission (%)	Filter designation
10771.5	31	79	a
10832.5	25	81	b
10893.0	28	78	c

Fig. 1. Percentage of filter transmission of the $\psi(25)$ photometric system.

The probable error of a single observation of the $\psi(25)$ index is 0.004 mag. Zirin's data (1982) have a threshold of detectability around 30 mA which is less accurate by a factor of two, roughly, than our photometric data. However, it should be pointed out that the presence of $\lambda 10832 \text{ \AA}$ water vapor line could be a cumbersome problem in photometric work. The amount of water vapor at San Pedro Observatory is between 1 and 3 mm precipitable water, most of the time (Gimmestad et al. 1972), fortunately.

The new photometric data, in magnitudes, are listed in Table 2.

III. DISCUSSION

The data contained in Table 2 comprise 31 stars which according to Zirin (1982) have all a zero equivalent width. Thus it seems reasonable to use these values to determine the constant, k , in equation (1). We obtain, after weighting the quantities by the total number of observations,

$$k = 0.167 \pm 0.005 \quad (2)$$

Actually the values of $\psi(25)$ given in Table 2 have been obtained applying equations (1) and (2). This means that, in the first approximation, if $\psi(25) = 0$, then $W(\text{mA})$ vanishes, probably. However, it should be noted that the great majority of the stars earlier than F0 so far observed have a negative $\psi(25)$ index, while the great majority of the stars listed in Table 2 from F0 to M7 have a positive $\psi(25)$ index.

At present the relationship between $\psi(25)$ and $W(\text{mA})$ is not very useful because more than half of the equivalent widths of our stars are zero, and also because a significant number of the remaining stars have a variable equivalent width (Zirin 1982), and a variable $\psi(25)$ index (see Table 2).

Between Zirin's results (1982) and the data contained in Table 2, we have 14 stars with a probable $\lambda 10830 \text{ \AA}$ variable in intensity, and 12 stars suspected of variability. Thus the large number of stars with a variable $\lambda 10830 \text{ \AA}$ line found by Zirin (1982) is confirmed. It is interesting to mention that the largest variation of this helium line, in our data corresponds to Algol. This large value was also found by Zirin.

The photometric data are too scanty to deduce at this stage photometric properties. However, it is interesting to note that two groups of stars, namely, the F-G type stars of luminosity class I, and the A-type stars of lower luminosity (III-V), all nearly lie in a straight line in the $\alpha(16)\Lambda(25)$ plane as shown in Figure 2. This relationship indicates that

TABLE 2

He I ($\lambda 10830$ A) Line Photometry of Bright Stars
(in magnitudes)

BS	Name	ψ (25)	n	MK	Rks	BS	Name	ψ (25)	n	MK	Rks
15	α And	-0.014	3	B8 IVp MnHg		5056	α Vir	-0.012	5	B1 III-IV + B2 V	
21	β Cas	0.005	5	F2 III-IV		5287	π Hya	-0.002	2	K2 IIIa-IIIb CN-0.5	
39	γ Peg	-0.006	5	B2 IV		5340	α Boo	0.007	5	K1 IIIb CN-1	
168	α Cas	0.003	5	K0 III	v:(2)	5429	ρ Boo	0.004	2	K3 II	
188	β Cet	0.022	3	K0 III CH-1H,K-0.5	v(2)	5506	ϵ Boo	0.005	2	K0 II-III	v(1)
264	γ Cas	-0.262	5	B0.5 IVe		5603	σ Lib	0.002	2	M3 IIIa	
337	β And	0.002	4	M0 IIIa		5744	ι Dra	0.012	2	K2 III	v:(1)
403	δ Cas	-0.011	5	A5 III-IVv		5854	α Ser	0.006	5	K2 IIIb CN-1 Fe 4143-1	
553	β Ari	-0.012	4	A5 V		6056	δ Oph	0.007	2	M0.5 III	
603	γ^1 And	-0.002	4	K3 I Ib	v(2)	6134	α Sco	-0.010	4	M1.5 Iab-Ib + B4 Ve	
617	α Ari	0.005	4	K2 IIIab Ca-1		6299	κ Oph	-0.002	2	K2 III	v:(1)
622	β Tri	-0.013	4	A5 III		6418	π Her	0.005	2	K3 IIab	
681	θ Cet	-0.020	3	M7 IIIe		6536	β Dra	0.037	2	G2 Ib-IIIa	v(2)
911	α Cet	0.007	4	M1.5 IIIa		6556	α Oph	-0.002	2	A5 III	
915	γ Per	0.003	4	G8 III + A2 V		6603	β Oph	0.009	2	K2 III	v:(2)
936	β Per	0.013	4	B8 V	v(1,2)	6859	δ Sgr	0.001	1	K3 IIIa	
1017	α Per	0.007	5	F5 Ib		7001	α Lyr	-0.021	7	A0 Va	
1122	δ Per	0.003	5	B5 III	v:(1)	7121	σ Sgr	0.016	1	B2.5 V	
1165	η Tau	-0.005	4	B7 IIIe		7194	ζ Sgr	-0.019	1	A2 III + A4 IV	
1203	ζ Per	-0.006	3	B1 Ib	v:(1)	7234	τ Sgr	0.002	1	K1 III	
1220	ϵ Per	-0.006	4	B0.5 V + A2 V		7235	ζ Aql	-0.017	1	A0 Vn	
1231	γ Eri	0.005	3	M0.5 III Ca-I Cr-I		7264	π Sgr	0.013	1	F2 II	
1457	α Tau	0.003	6	K5 III		7310	δ Dra	0.009	2	G9 III	
1577	ι Aur	-0.006	3	K3 II	v(2)	7417	β Cyg	0.004	2	K3 II + B0.5 V	v(2)
1605	ϵ Aur	0.001	4	F0 Iae		7525	γ Aql	0.025	2	K3 II	v(2)
1666	β Eri	-0.010	2	A3 III		7528	δ Cyg	-0.014	2	B9.5 IV + F1 V	v:(1)
1708	α Aur	0.032	4	G5 IIIe + G0 III	v:(2)	7557	α Aql	-0.004	8	A7 V	
1713	β Ori	0.039	2	B8 Iae	v(2)	7615	η Cyg	0.005	4	K0 III	v(1)
1790	γ Ori	-0.001	4	B2 III		7635	γ Sge	0.002	1	M0 III	
1791	β Tau	-0.007	4	B7 III		7751	θ^2 Cyg	0.012	2	K3 Ib + B3 V	
1829	β Lep	0.011	2	G5 II		7776	β Cap	0.015	1	F8 V + A0	
1852	δ Ori	-0.012	3	B0 III + O9 V		7796	γ Cyg	0.014	3	F8 Ia	
1865	α Lep	0.004	2	F0 Ib		7924	α Cyg	0.000	8	A2 Iae	
1899	ι Ori	-0.006	2	O9 III	v(2)	7949	ϵ Cyg	0.008	7	K0 III	v:(1)
1903	ϵ Ori	-0.039	2	B0 Iae	v(1)	8162	α Cep	-0.003	1	A7 V	
1910	ζ Tau	0.003	4	B4 IIIpe		8232	β Aqr	0.021	3	G0 Ib	
1948	ζ Ori	-0.040	2	O9.5 Ibe	v:(1)	8308	ϵ Peg	-0.004	2	K2 Ib	v(2)
1956	α Col	-0.011	1	B7 IVe		8322	δ Cap	-0.001	2	Amv	
2004	κ Ori	-0.034	1	B0.5 Iav		8353	γ Gru	-0.008	2	B8 III	v:(1)
2061	α Ori	-0.008	3	M1-2 Ia-Iab	v:(2)	8414	α Aqr	0.024	2	G2 Ib	v(2)
2088	β Vir	-0.019	3	A2 IV		8650	η Peg	0.005	2	G2 II-III + F0 V	
2095	θ Aur	-0.012	2	A0p Si		8728	α PsA	-0.018	1	A3 V	
2286	μ Gem	0.006	1	M3 IIIab		8775	β Peg	0.003	3	M2.5 II-III	
2421	γ Gem	-0.012	1	A0 IV		8781	α Peg	-0.013	5	B9 V	
4932	ϵ Vir	0.006	2	G8 IIIab							

Remarks to Table 2: v, variable; v: suspected of variability; (1) this photometry; (2) Zirin (1982).

for these pair of stellar groups, the stronger the absorption in the Balmer hydrogen $H\alpha$ line, the weaker the absorption in the neutral helium line at $\lambda 10830$ A. Perhaps this is extended to weak emission (in He I) for the strongest absorption (in $H\alpha$).

Actually the F,G-type stars plotted in Figure 2 have a positive $\psi(25)$, while the A-Type stars also plotted in Figure 2 have $\psi(25)$ index from nearly zero, but negative, to negative values. For these stars, as expected, there is not a linear relationship between $\Lambda(9)$ and $\psi(25)$ photometric indices as shown in Figure 3.




Fig. 2. The $\alpha(16)\psi(25)$ plane. Coding: filled circles, A-type stars of low luminosity (m, Am; n, An; p, Ap; sb, A+A -see Table 2); crosses, F, G-type stars of luminosity class Ib; crossed circle, luminosity class Ia (F8); crossed square, luminosity class Ib-II (G2).

Fig. 3. The $\Lambda(9)\psi(25)$ plane. Coding as in Figure 2.

IV. CONCLUSION

The photometric technique to measure total absorptions of spectral lines has been very fruitful (cf. Strömberg 1963; Mendoza loc. cit.). This preliminary work indicates that for a given telescope, the limit magnitude reached to measure the $\lambda 10830$ Å line is fainter, when one uses photometric indices, than that of spectroscopic techniques. Unfortunately, no line profile is obtained through the former method.

We hope to observe this neutral helium line next year with Silicon detectors which are more sensitive than the photomultiplier employed in this preliminary work.

ACKNOWLEDGEMENTS

We are indebted to Mr. E.T. Mendoza for his invaluable help during the 1986 observations. We also thank Mrs. D. Learn for suggesting changes in the wording, and Mr. A. García for inking the figures.

REFERENCES

- Johnson, H.L. and Mitchel, R.I. 1975, *Rev. Mexicana Astron. Astrof.*, 1, 299.
 Gimmetstad, G.G., Gebbie, H.A., Bohlander, R.A., and Mendoza, E.E. 1972, *Ap. J.*, 178, 267.
 Mendoza, E.E. 1973, *IAU Symposium No. 50, Spectral Classification and Multicolor Photometry*, ed. Ch. Ferenbach and B.E. Westerlund, (D. Reidel Publishing Co. Dordrecht-Holland), p. 208.
 Mendoza, E.E. 1975, *Pub. Astron. Soc. Pacific* 87, 505.
 Mendoza, E.E. 1976, *IAU Symposium No. 72, Abundances Effects in Classification*, ed. B. Hauck and B.C. Keenan, (D. Reidel Publishing Co. Dordrecht-Holland), p. 364.
 Mendoza, E.E. 1979a, *IAU First Latin American Regional Astronomy Meeting*, ed. A. Gutiérrez-M and H. Moreno (Publ. Dept. Astron. Univ. Chile), p: 250.
 Mendoza, E.E. 1979b, *Astron. Astrophys.*, 71, 147.

- Mendoza, E.E. 1981, *IAU Symposium No. 98, Be Stars*, ed. M. Jaschek and H.G. Groth, (D. Reidel Publishing Co. Dordrecht-Holland), p. 3.
- Mendoza, E.E. 1983, *IAU Standard Star Newsletter*, ed. L.E. Pasinetti, No. 2, p. 12.
- Mendoza, E.E. 1986, *IAU Colloquium No. 92, Physics of Be Stars* (in press).
- Strömberg, B. 1963, *Stars and Stellar Systems*, ed. G.P. Kuiper and B.M. Middlehurst (Chicago: The University of Chicago Press), Vol. 3, Chap. 9, p. 123.
- Zirin, H. 1982, *Ap. J.*, 260, 655.

E.E. Mendoza: P.O. Box 20-158, México 20, D.F., 01000 México.