

## RADIO MAP OF THE REGION G331.5-0.1 AT 22 GHz

Luiz H. Amaral and Zulema Abraham

Instituto Astronômico e Geofísico, USP

RESUMEN. Se hicieron relevamientos en 22 GHz en la región térmica G331.5-0.1 con el radiotelescopio de Itapetinga, el cual tiene una resolución de 4.6 minarc en esta frecuencia. La región incluye también las fuentes de radio G331.5-0.1, G331.4-0.0, G331.3-0.2, G331.3-0.3, G331.0-0.1, G331.1-0.5, G330.9-0.4 y G330.7-0.4, conocidas en 5 GHz.

ABSTRACT. The thermal region G331.5-0.1 was mapped at 22 GHz with the Itapetinga radiotelescope, which has a resolution of 4.6 arcmin at this frequency. The region includes also the radio sources G331.5-0.1, G331.4-0.0, G331.3-0.2, G331.3-0.3, G331.0-0.1, G331.1-0.5, G330.9-0.4, and G330.7-0.4, known at 5 GHz.

Key words: RADIO SOURCES-GENERAL

## . INTRODUCTION

We present a 22 GHz radio map of the regions of the galactic plane centered in  $6^{\text{h}}7^{\text{m}}30^{\text{s}}$  of right ascension and  $-51^{\circ}39'24''$  of declination. This region is located in the direction of the Norma spiral arm and presents 8 sources, the most intense, G331.5-0.1 gives the name to it. Maps at 5 GHz with 4' resolution and at 408 MHz with a 3' resolution are available (Haynes et al. 1978, Goss e Shaver 1970). Recombination lines and strong H<sub>2</sub>O maser emission are also characteristic of some of the sources in this region.

## I. OBSERVATIONS

The observations were made with the 13.7 m Itapetinga radiotelescope at the frequency of 22 GHz. The receiver was a K-band mixer with a d.s.b. of 1 GHz. The receiver was operated in the total power mode, the system temperature was around 700 K and the integration time 50ms. The observations consisted of scans in right ascension (RA) with an amplitude of  $^{\circ}$ , separated in declination by 2'. During each observation there were made 30 scans of 20 s duration, preceded by calibrations with a noise source of known temperature and a load at room temperature. Before each observing period, measurements of Virgo A were made, in the form of scans in azimuth and elevation, to determine the antenna efficiency and verify the pointing accuracy.

## II. RESULTS

In Fig. 1 we present the 22 GHz map of the G331.5-0.1 region with 4.6 resolution. The numbers (N) associated with each contour line are related to the antenna temperature  $T_A$  by  $T_A = 0.359 (N/100)$ . The relationship between antenna temperature and flux density ( $S_p$ ) for a point source was obtained from the observations of Virgo A, which has a flux density of 1.5 Jy at 22 GHz (Janssen et al. 1974). We found the relation  $S_p(\text{Jy}) = 63.2 \times T_A(\text{k})$ .

We corrected the antenna temperature for the effects of atmospheric absorption, using the calibration method described by Abraham (1986).

The physical parameters of the sources detected in the map are presented in table 1. Columns 1, 2 and 3 correspond to the name and coordinates (1950) of the sources, columns 4 to 8 represent respectively, the peak antenna temperature, peak flux density, angular size, integrated flux density and spectral index between 5 and 22 GHz. The thermal (T) or non-thermal (NT) nature of the sources is indicated in column 9. The magnitude of

the spectral index was calculated using the 5 GHz data of Caswell et al. (1987). The detection of the hydrogen recombination lines H109 $\alpha$  and H110 $\alpha$  by these authors and the determination of their physical parameters confirm the thermal nature of the sources. Due to its low intensity we were not able to determine the size of the source G330.7-0.4.




Figure 1 - Radio map of the region G331.5-0.1 at 22 GHz

TABLE 1

FONTE	A.R. (1950)			DEC (1950)			$T_A$ (K)	$S_p$ (Jy)	$\theta$ (')	$S_I$ (Jy)	$\alpha$	NAT
	h	m	s	o	"	'						
G331.5-0.1	16	08	21.6	-51	19	24	0.359	22.7	3.2	33.5	0.11	T
G331.4-0.0	16	07	17.2	-51	23	24	0.084	5.3	3.1	7.7	0.17	T
G331.3-0.3	16	08	34.5	-51	39	24	0.090	5.7	2.5	7.4	0.03	T
G331.3-0.2	16	07	36.2	-51	35	24	0.067	4.2	2.2	5.1	0.12	T
G331.0-0.1	16	06	12.4	-51	47	24	0.095	6.0	7.5	21.7	0.13	T
G331.1-0.5	16	08	21.9	-51	55	24	0.070	4.4	2.1	5.4	-0.21	T
G330.9-0.4	16	06	25.1	-51	59	24	0.114	7.2	2.7	9.7	0.06	T
G330.7-0.4	16	05	45.8	-52	07	24	0.038	2.4	---	---	---	-

## ACKNOWLEDGEMENTS

We acknowledge the financial support of the Brazilian agencies FAPESP and CNPq.

## REFERENCES

- Abraham, Z.; Botti, L.C.L.; del Ciampo, L.F., 1986, *Rev. Mexicana Astron. e Astrof.* 12, 414  
Caswell, J.L.; Haynes, R.F., 1987, *Astron. Astrophys.* 171, 261  
Haynes, R.F.; Caswell, J.L.; Simons, L.W., 1978, *Aust. J. Phys. Astrophys. Suppl.* no. 45, 1  
Janssen, M.A.; Golden, L.M.; Welch, W.J., 1974, *Astron. Astrophys.* 33, 373  
Shaver, P.A.; Goss, W.M., 1970, *Aust. J. Phys. Astrophys. Suppl.* 14, 1-133

Luiz H. Amaral and Zulema Abraham: Instituto Astronômico e Geofísico, Universidade de São Paulo,  
Caixa Postal 30627, CEP 01051, São Paulo, SP, Brazil

