

star infrared color box. In the present form we search for optical objects within the boundaries defined by Weintraub (1990, ApJS, 74, 575). There are about 3500 of such sources in both hemispheres, 20% having possible optical counterparts brighter than 14.5 mag, as a result of the correlation with the Guide Star Catalogue (GSC).

We have taken coudé spectra with CCD in the 6500 Å region, and we defined as a T Tauri star (TTS) an object having both the Li absorption line and H α emission line. We found 45 sources associated with new TTS, 7 being optical pairs and for 28 we suspect they are TTS. We also suspect the Herbig Ae/Be nature for 76 objects. In two cases the object may be a Fuori-like star. We found 6 new late type Li-rich giant stars. Some of the new found YSO are high latitude objects. The survey is now 80% complete south of +30°.

MOLECULAR GAS IN CENTAURUS A. THE ^{12}CO J = 2-1 MAP

W. Wild¹, M. Cameron², A. Eckart², R. Genzel²,
H. Rothermel², G. Rydbeck³, and T. Wiklind³

We mapped the ^{12}CO J = 2-1 line in Centaurus A using the 15-m Swedish-ESO Submillimetre Telescope (SEST) on La Silla, Chile. Centaurus A (NGC 5128) is a nearby (3 Mpc) elliptical galaxy with a prominent dust lane, extensive radio lobes and a compact radio continuum source. Due to its proximity and peculiar morphology, it has been observed over a large part of the electromagnetic spectrum. The detailed study of the molecular interstellar medium, however, has begun only recently with the availability of a large millimeter telescope in the southern hemisphere.

We present a map of the ^{12}CO J = 2-1 emission along the dust lane of Centaurus A. In several observing runs between December 1990 and May 1992 we measured a total of 240 positions extending over an area of $200'' \times 70''$. The angular resolution of SEST at the frequency of the ^{12}CO J = 2-1 transition (230.5 GHz) is $22''$. The grid spacing was typically $8''$ in the inner parts of the dust lane and $16''$ in the outer parts. Integration times per position varied between 4 and 30 minutes.

Strong emission in the ^{12}CO J = 2-1 line (up to a level of $T_{mb} = 0.6$ K) is seen over a large part of the dust lane. The emission is generally symmetrical about the nucleus but, as in the case of the J = 1-0 and 50 micron maps, it is not centrally peaked. The striking similarity in the morphologies of the ^{12}CO J = 2-1, J = 1-0 and 50 micron maps suggests that the gas and warm dust are probably well coupled. The good spatial sampling of our map has allowed

us to investigate into the kinematics of the molecular gas (Rydbeck et al. 1993, in preparation). The excitation conditions in the disk can be probed using the ^{12}CO J = 2-1/1-0 ratio, when the J = 2-1 map has been convolved to the resolution of the 1-0 map. This results in a ratio of close to unity at the position of the nucleus, a value which is also typical of the gas throughout the whole extent of the disk. Such a high J = 2-0/1-0 ratio implies that the bulk of the gas in the dust lane is warm ($T > 15$ K), dense ($n(\text{H}_2) 2 \times 10^4 \beta \text{ cm}^{-3}$, the critical density required to thermalise the J = 2-1 level, where β is the escape probability), and, probably partially, optically thick. This conclusion is supported by a ^{13}CO J = 2-1/1-0 ratio of 0.9 at one position in the disk.

¹ European Southern Observatory, La Silla, Chile.

² Max-Planck-Institut für extraterrestrische Physik, Germany.

³ Onsala Space Observatory, Sweden.

BLUE STRAGGLERS IN OPEN CLUSTERS

Javier Ahumada and Emilio Lapasset

Observatorio Astronómico

Universidad Nacional de Córdoba, Argentina

A statistical study of the observational characteristics of blue stragglers in open clusters is presented. It is based on the visual inspection of color-magnitude diagrams of the clusters with photometry published before December 1991. According to membership probability and quality of the observations, the blue stragglers have been classified into three categories. Some interesting relations as: number of blue stragglers against cluster ages, number of blue stragglers versus number of ordinary stars per cluster, and degree of concentration of the blue stragglers in each cluster are shown.

OPTICAL IDENTIFICATION OF ROSAT X-RAY SOURCES AT THE GUILLERMO HARO OBSERVATORY

I. Appenzeller¹, C. Chavarría², L. Corral², T. Fleming³,
G. Hasinger³, J. Krautter¹, R. Mújica², M. Pakull¹,
L. Terranegra², J. Trümper³, and F.J. Zickgraf¹

First results of an optical identification program of X-ray sources newly discovered during the ROSAT All-Sky Survey are presented. The sample comprises about 1300 sources which are contained in six "study areas" of about 150 square degrees each. Selection criteria for these areas were: (i)