

# A METHOD OF NUMERICAL INTEGRATION FOR CALCULATING ASTRONOMICAL REFRACTION

J. Stock and R. Molina

Centro de Investigaciones de Astronomía, CIDA, Venezuela

*Received 1998 January 26; accepted 1998 July 16*

## RESUMEN

Se calcula la refracción astronómica basada en una integración numérica de ecuaciones diferenciales, las cuales describen la propagación de un rayo de luz a través de una atmósfera estratificada de manera concéntrica. El cálculo hace uso de una atmósfera patrón y de las condiciones meteorológicas locales en el Observatorio Nacional de Llano del Hato en Mérida, Venezuela ubicado a 3600 m de altura. Se deriva una fórmula de interpolación la cual permite un cálculo rápido de la refracción. La comparación de éste, con otros métodos, arroja diferencias mínimas.

## ABSTRACT

Astronomical refraction is calculated on the basis of a numerical integration of differential equations which describe the propagation of a beam of light through a concentrically stratified atmosphere. The calculations make use of the standard atmosphere and the local meteorological conditions at the National Observatory Llano del Hato in Mérida, Venezuela located at an elevation of 3600 m. An interpolation formula is derived which permits a rapid calculation of the refraction, and when compared with other methods shows only minor differences.

*Key words:* **ASTROMETRY — ATMOSPHERIC EFFECTS**

## 1. INTRODUCTION

‘Astronomical refraction’, i.e., the correction which has to be applied to a measured zenith distance in order to eliminate the effect of the deviation of a light beam as it passes through the atmosphere, has been the subject of many studies. The exact determination of this correction is only possible through a comparison of the measured angles with an exact and error free system of astronomical coordinates. A number of reasons, mostly of instrumental nature, and uncertainties in the refraction correction, have so far impeded the construction of such a ‘reference system’. The Hipparcos catalogue will make such a comparison possible and thus will tell us which of the methods developed so far comes closest to the true refraction.

The refraction of light in the air is strictly a function of the gradient of the refractive index, and the latter is a function of the density of the air. Thus, if we had an exact model of the atmosphere, i.e., an exact description of the density as a function of

the three-dimensional space coordinates  $X$ ,  $Y$ , and  $Z$ , we can follow a light beam through the air, as proposed for geodetic applications by Stock (1986). This is done with a step by step numerical integration of a set of differential equations and thus differs from the methods used by other authors; for example, Fukawa & Yoshizawa (1985), Yatsenco (1995), or Stone (1996). The latter states that ‘*atmospheric refraction should be determined ideally by tracing the path of light through the Earth’s atmosphere*’. This is exactly what this paper pretends to do.

## 2. GEOMETRY OF THE LIGHT PATH

We shall make the assumption that the structure of the atmosphere is described by concentric shells; hence the refractive index  $\eta(h)$  is only a function of the altitude  $h$  above sea level. The geometry of the light path is shown in Figure 1, where the observed zenith distance is  $z_0$ ;  $z$  is the angle of the beam with respect to the  $Y$ -axis at the point  $X, Y$ ;  $z'$  is the angle of the beam with respect to the local zenith at  $X, Y$ ;




Fig. 1. Geometry of a light beam as it passes through a concentric atmosphere. Symbols are explained in the text.

$R_0$  is the radius of the Earth at the observing site, and  $\theta$  is the angle relative to the at the center of the Earth.

Let  $s$  be a coordinate along the light path. Then, according to Stock (1986), the variation of  $z'$  as the beam passes from  $s$  to  $s + ds$  is given by

$$\frac{dz'}{ds} = -\frac{1}{\eta(h)} \sin(z') \frac{d\eta(h)}{dh}. \quad (1)$$

This equation follows from differentiating Snell's law of refraction and then applying it to the light beam in Fig. 1. Furthermore, we have the following relations

$$\frac{dx}{ds} = \cos(z), \quad (2)$$

and

$$\frac{dy}{ds} = \sin(z). \quad (3)$$

Evidently, with a knowledge of  $d\eta(h)/dh$  this system of differential equations can be solved.

### 3. THE ATMOSPHERIC MODEL

To construct the function  $\eta(h)$  we use the physical data of the Earth's atmosphere of the Standard Atmosphere (1976), which gives the physical parameters of the air at each elevation. These, together with the measured meteorological parameters at the observing site, are interpolated by exponential functions, thus constructing continuous functions which extend to an elevation of 90 km, where the mentioned

standard data end. From these functions the density  $\rho$  of the air can be calculated for any elevation with the help of known relations. The latter, for instance, may be found in § 3 of the paper by Stone (1996). The refractive index is then calculated from

$$\eta(h, \lambda) = a(\lambda)\rho(h), \quad (4)$$

where  $\lambda$  is the wavelength, and  $a(\lambda)$  a well known function.

### 4. RESULTS

The above integration was carried out for a specific site, namely the Astronomical Observatory in the Venezuelan Andes, located at an elevation of 3600 m. Our results refer to the visual wavelengths, 5500 Å, and calculations range from 0° to 80° zenith distance. For the purpose of interpolation and a more rapid handling of the results a formula of the form

$$R = A \tan(z) + B \tan^3(z), \quad (5)$$

was fitted to the data by least squares. The results of the polynomial are given in the Table 1. The coefficients  $A$  and  $B$  are given at the bottom of the table.

Of the standard atmosphere we need the density as function of the elevation  $h$  which can readily be converted into the respective refractive index. The integration is started at an elevation of 3600 m for which the density and hence, the refractive index, is calculated from the average local night-time meteorological parameters. The next point is taken from the table of the Standard Atmosphere at an elevation of 4000 m. We should mention here that the value calculated for the initial point agrees closely with what would be obtained directly from the table. From then on, points of the relation of the refractive index versus the elevation were taken from the table at increasing intervals, from 1000 m in the lower troposphere to 20,000 m at the top of the stratosphere. The refractive index was interpolated from point to point by an exponential function. The gradient of the refractive index, which is needed in equation (1), was calculated numerically from the interpolating functions. The refractive index is in this manner represented by a continuous function. Its derivative is discontinuous at the junction points, but the numerically calculated differences have no discontinuity. The path length of the integration was chosen as 5.0 m up to an elevation of 10,000 m, and then increased up to 50.0 m at the end of the integration at an elevation of 120 km. The results are given in Table 1.

At this point, a comparison with other methods and at different wavelengths may be of interest. Comparisons can be made with the widely used Pulkovo Table (Orlov 1956) and with data by Fukawa & Yoshizawa (1985) for 5753 Å. The differences be-

TABLE 1

ASTRONOMICAL REFRACTION AT THE NATIONAL OBSERVATORY  
LLANO DEL HATO IN MERIDA, VENEZUELA

Zenithal Distance $z_0(^{\circ})$	Atmospheric Refraction $R_0(^{\prime\prime})$	$A \tan(z)$ $+B \tan^3(z)^a$ $R_0(^{\prime\prime})$	Zenithal Distance $z_0(^{\circ})$	Atmospheric Refraction $R_0(^{\prime\prime})$	$A \tan(z)$ $+B \tan^3(z)^a$ $R_0(^{\prime\prime})$
1 .....	0.690	0.689	41 .....	34.319	34.306
2 .....	1.380	1.379	42 .....	35.543	35.532
3 .....	2.071	2.070	43 .....	36.811	36.797
4 .....	2.763	2.762	44 .....	38.117	38.103
5 .....	3.457	3.456	45 .....	39.469	39.454
6 .....	4.153	4.152	46 .....	40.867	40.852
7 .....	4.851	4.850	47 .....	42.317	42.302
8 .....	5.553	5.551	48 .....	43.822	43.806
9 .....	6.257	6.256	49 .....	45.387	45.369
10 .....	6.966	6.965	50 .....	47.014	46.996
11 .....	7.679	7.678	51 .....	48.708	48.692
12 .....	8.398	8.395	52 .....	50.478	50.461
13 .....	9.121	9.119	53 .....	52.328	52.311
14 .....	9.851	9.848	54 .....	54.263	54.247
15 .....	10.586	10.583	55 .....	56.295	56.278
16 .....	11.329	11.325	56 .....	58.429	58.412
17 .....	12.079	12.075	57 .....	60.676	60.657
18 .....	12.837	12.833	58 .....	63.041	63.023
19 .....	13.603	13.599	59 .....	65.547	65.528
20 .....	14.378	14.374	60 .....	68.195	68.178
21 .....	15.165	15.160	61 .....	71.007	70.992
22 .....	15.961	15.956	62 .....	74.000	73.985
23 .....	16.768	16.763	63 .....	77.191	77.178
24 .....	17.587	17.582	64 .....	80.606	80.594
25 .....	18.421	18.414	65 .....	84.270	84.258
26 .....	19.267	19.260	66 .....	88.211	88.202
27 .....	20.128	20.120	67 .....	92.465	92.460
28 .....	21.003	20.995	68 .....	97.075	97.075
29 .....	21.894	21.887	69 .....	102.094	102.096
30 .....	22.804	22.796	70 .....	107.575	107.582
31 .....	23.731	23.724	71 .....	113.588	113.603
32 .....	24.680	24.671	72 .....	120.220	120.244
33 .....	25.648	25.639	73 .....	127.578	127.609
34 .....	26.639	26.629	74 .....	135.786	135.827
35 .....	27.652	27.642	75 .....	145.010	145.057
36 .....	28.690	28.680	76 .....	155.445	155.500
37 .....	29.756	29.745	77 .....	167.351	167.410
38 .....	30.848	30.839	78 .....	181.071	181.116
39 .....	31.973	31.962	79 .....	197.039	197.046
40 .....	33.127	33.117	80 .....	215.861	215.763

<sup>a</sup>  $A = 39.5006'' \pm 0.0021''$ ;  $B = -0.0450'' \pm 0.0001''$ .

tween the data from these two sources and ours are shown in Figure 2. Comparisons at two other wavelengths can also be made with the data from Fukawa

& Yoshizawa. The respective results are shown in Figure 3. From the figures, one can conclude that the different methods of calculating the astronomical




Fig. 2. The refraction of this paper minus the refraction from the Pulkovo Table and the calculations by Fukawa & Yoshizawa as function of the zenith distance for 5753 Å.




Fig. 3. The refraction of this paper minus the refraction calculated by Fukawa & Yoshizawa for 5500 Å and 5936 Å as function of the zenith distance.

refraction lead to practically identical results down to a zenith distance of 70°. Beyond this distance a major discrepancy builds up.

## 5. CONCLUSIONS

We have shown that accurate information on astronomical refraction can be obtained by directly integrating a light beam. The results are very close to those obtained by other methods. In fact, for zenith distances < 78° the differences are less than 200 mas. Comparison with other independent data is needed, in order to determine which method comes closest to the true refraction. The Hipparcos catalogue may provide the necessary reference frame.

## REFERENCES

- Fukawa, R., & Yoshizawa, M. 1985, PASJ, 37, 747
- Orlov, B. A. 1956, Refraction Tables of the Pulkovo Observatory, 4th edition (Moscow: Academic of Sciences)
- Standard Atmosphere. 1976, National Oceanic and Atmospheric Administration, NASA, and USAF, NOAA-S/T76-1562 (Washington, D. C.: U. S. Government Printing Office)
- Stock, J. 1986, Tectonophysics, 130, 179
- Stone, R. C. 1996, PASP, 108, 1051
- Yatsenko, A. Y. 1995, A&AS, 111, 579