

DETAILED ATMOSPHERIC ABUNDANCE ANALYSIS OF THE OPTICAL COUNTERPART OF THE IR SOURCE IRAS 16559-2957

R. E. Molina¹ and A. Arellano Ferro²

Received 2012 June 8; accepted 2013 January 16

RESUMEN

Hemos emprendido un análisis detallado de las abundancias químicas de la contraparte óptica de la fuente infrarroja IRAS 16559-2957 con el fin de confirmar su posible naturaleza de estrella post-AGB. El análisis de un gran número de elementos, incluidos CNO y ¹²C/¹³C, muestra que este objeto ha experimentado el primer dragado y parece estar aún en fase de RGB.

ABSTRACT

We have undertaken a detailed abundance analysis of the optical counterpart of the IR source IRAS16559-2957 with the aim of confirming its possible post-AGB nature. The star shows solar metallicity and our investigation of a large number of elements including CNO and ¹²C/¹³C suggests that this object has experienced the first dredge-up and it is likely still at RGB stage.

Key Words: stars: abundances — stars: AGB and post-AGB — stars: evolution

1. INTRODUCTION

Post-AGB stars (PAGB's) are at a very advanced stage of evolution. They show peculiarities in the observed elements in their atmospheres (e.g., CNO and s-process elements abundances) which are connected with the dredge-up episodes that occur as the star evolves up on the AGB. PAGB's also have a high mass loss ($\geq 10^{-6} M_{\odot} \text{ yr}^{-1}$) which obscures the central star and produces emission in the IR and OH masers (Zijlstra et al. 1989). Consequently, the IRAS colour-colour diagram ([25]–[60],[12]–[25]) and its IRAS variability has been widely used in the identification of PAGB and proto-planetary nebulae (PPN) candidates (van der Veen & Habing 1988; García-Lario et al. 1997; Suárez et al. 2006; Szczerba et al. 2012).

The identification of optical counterparts of IR and OH masers and IRAS sources is of fundamental importance since it is in the optical spectrum that the abundance anomalies of structural and evolutionary relevance can be detected and studied, such as CNO abundances and those of s-process elements (e.g., La, Ba, Nd, Sr, Y, Zr).

The IRAS 16559-2957 source has been included in several studies aimed to identify and evaluate the evolutionary status of large lists of PAGB and PPN candidates. In the finding chart provided by Hu et al. (1993) (their object 34) there is only one bright star within the IRAS beam. According to Hu et al. (1993) it is a F5I star of $V \sim 13$ with H α filled in. The presence of a circumstellar envelope around a central star was confirmed by the detection of strong OH maser at 1612 and 1665 MHz (Hu et al. 1993). The radial velocity of the OH maser profile varies between 57–70 km s⁻¹ (te Lintel Hekkert et al. 1991).

The source was included in the sample of PAGB and PN candidates studied by Suárez et al. (2006) who find no optical counterpart. However, Ramos-Larios et al. (2009), in their search of heavily obscured PAGB and PN stars, found a near-IR counterpart in the 2MASS Point Source Catalogue and an optical counterpart in the Digital Sky Survey, which corresponds to a $V \sim 13.2$ star (see their finding chart on Figure 3), in agreement with Hu et al. (1993). Then the star has been *bona fide* catalogued as a PAGB (e.g., Szczerba et al. 2007; SIMBAD³ data base).

With the aim of confirming the PAGB nature of the optical counterpart we have studied a high resolution optical spectrum of the $V \sim 13.2$ star at

¹Laboratorio de Investigación en Física Aplicada y Computacional, Universidad Nacional Experimental del Táchira, Venezuela.

²Instituto de Astronomía, Universidad Nacional Autónoma de México, Mexico.

³<http://simbad.u-strasbg.fr/simbad/sim-fid>.

$\alpha(2000) = 16^{\text{h}}59^{\text{m}}08.2^{\text{s}}$, $\delta(2000) = -30^{\circ}01'40.3''$. In this paper we present a detailed photospheric chemical abundances analysis of this optical source and we shall refer to the object as IRAS 16559-2957.

The paper is organized as follows: in § 2 the observations and reductions are described, in § 3 we discuss the adopted atmospheric parameters and the reddening, in § 4 we employ the equivalent widths and spectral synthesis approaches to derive the photospheric chemical abundances, in § 5 we discuss our results, and in § 6 we summarize our conclusions.

2. OBSERVATIONS AND REDUCTIONS

The spectrum of IRAS 16559-2957 ($V = 13.2$) was obtained on February 27, 2008 with the 2D Coudé echelle spectrograph (Tull et al. 1995) on the 2.7 m telescope at the McDonald Observatory giving $\sim 40,000$ resolution and a wavelength coverage from 3900 to 10000 Å. The spectrum was reduced using the IRAF astronomical routines. The equivalent widths (EW's) were measured using the SPLOT task and their accuracy is generally better than 10% for spectra with S/N ratio larger than 50. We generally restricted ourselves to unblended weak features and avoided using lines stronger than 200 mÅ.

We measure the heliocentric radial velocity using the Doppler shift of 40 clean unblended absorption lines that cover a wide spectral range. We derive a heliocentric radial velocity of $-2.3 \pm 0.5 \text{ km s}^{-1}$ ($V_{\text{LSR}} = +7.1 \text{ km s}^{-1}$).

3. ATMOSPHERIC PARAMETERS

Before we derive the elemental abundances it is necessary to estimate the fundamental atmospheric parameters; effective temperature, surface gravity and microturbulence velocity. We have approached the task as discussed in the following subsections.

3.1. Spectroscopy

An independent estimation of T_{eff} , $\log g$ and the microturbulence velocity ξ_t can be obtained from the behaviour of the lines of a well represented element (e.g., Fe), and the MOOG code (2009 version) by Sneden (1973), which is based on local thermodynamical equilibrium (LTE) calculations. We employed the atmospheric model collection of Castelli & Kurucz (2003).

We have therefore followed the standard procedure for temperature and gravity determinations. For the temperature, it is required that the Fe I abundances are independent of the low excitation potential of the lines. The gravity can be derived by requiring that the Fe I and Fe II lines yield




Fig. 1. The standard deviation of Fe I and Fe II abundances as a function of microturbulence velocity for IRAS 16559-2957. The dashed vertical line corresponds to the value of the adopted microturbulence velocity.

the same abundance. This procedure led to values $T_{\text{eff}} = 4250 \pm 200 \text{ K}$ and $\log g = 1.5 \pm 0.25$. This effective temperature does not correspond to a star of spectral type F5I but it is more typical of an early K-type star.

The microturbulence is estimated by requiring that the derived abundances are independent of the line strengths for a given species. Generally, lines of Fe II are preferred. It is known that Fe II lines are not seriously affected by departure from LTE (Schiller & Przybilla 2008). On the contrary Fe I lines are affected by NLTE effects (Boyarchuck, Lyubimkov, & Sakhbullin 1985; Thévenin & Idiart 1999). For cool stars, however, not many Fe II lines are available. The microturbulence velocity is therefore determined using the error diagram for Fe I or Fe II (depending on the number of measured lines). For a given model, we compute the dispersion in the Fe abundances over a range in the microturbulence velocity using the method of Sahin & Lambert (2009). Figure 1 shows the standard deviation σ_x as a function of microturbulence velocity ξ_t for Fe I and Fe II lines. From the minimum error we estimate $\xi_t = 1.43 \text{ km s}^{-1}$.

3.2. Determination of $E(B-V)$

The initial effective temperature of IRAS 16559-2957 could also be estimated from JHK photometry available in the work of Cutri et al. (2003) and the T_{eff} calibrations of Alonso, Arribas, & Martínez-Roger (1999) from the $J - H$ and $V - K$ colours.

TABLE 1

ATMOSPHERIC PARAMETERS
OF IRAS 16559-2957

T_{eff} (K)	$\log g$	ξ_t (km s ⁻¹)
4250	1.5	1.43

However, a previous knowledge of the reddening would be necessary. Instead we have used the calibrations to estimate the total extinction. The interstellar extinction maps of Schlegel, Finkbeiner, & Davis (1998) lead to a value $E(B - V) = 0.38$, and after applying the correction recommended by Bonifacio, Caffau, & Molaro (2000) for excesses larger than 0.15, we find $E(B - V) = 0.28$. Since the star is affected by circumstellar material responsible for the OH maser and IR excess, we have increased the value of $E(B - V)$ until the calibrations of Alonso et al. (1999) produced a value of $T_{\text{eff}} \sim 4250$ K, estimated spectroscopically. The total, interstellar plus circumstellar, reddening is $E(B - V) = 0.81$, leaving $E(B - V) = 0.53$ as our estimation of the circumstellar contribution.

The basic spectroscopic atmospheric parameters are summarized in Table 1.

The models of Castelli & Kurucz (2003) are calculated in steps of ± 250 K, ± 0.5 and ± 0.25 km s⁻¹ in T_{eff} , $\log g$ and ξ_t respectively. We have used however a set of interpolated models in steps of ± 100 K, ± 0.25 and ± 0.20 km s⁻¹. The sensitivity of the derived abundances to the uncertainties of atmospheric parameters is given in Table 2 and discussed in the next section.

4. ATMOSPHERIC ABUNDANCES

4.1. Equivalent Width Analysis

To calculate the atmospheric abundances we have used the 2009 version of the code MOOG (Snedden 1973) which performs LTE line analysis. We use the grid ATLAS9 of plane-parallel model atmosphere computed by Castelli & Kurucz (2003). Chemical abundances of well represented elements were determined through the analysis of equivalent widths using the *abfind* driver in MOOG code.

The oscillator strength or *gf* value is an important atomic parameter that affects the abundance calculations. The *gf* values can be obtained from numerous sources, and the uncertainties vary from element to element. For example, experimental values for Fe I and Fe II of high accuracy, between 5 to

TABLE 2

SENSITIVITY OF ABUNDANCES RELATIVE
TO THE UNCERTAINTIES IN THE MODEL
PARAMETERS FOR IRAS 16559-2957

Species	ΔT_{eff} +100 K	$\Delta \log g$ +0.25	$\Delta \xi_t$ +0.2 km s ⁻¹
C	+0.02	-0.05	0.00
N	-0.05	-0.08	0.00
O I	-0.02	-0.11	+0.01
Li I	-0.12	+0.05	0.00
Na I	-0.10	0.00	+0.10
Mg I	-0.01	-0.04	+0.09
Al I	-0.07	-0.03	+0.07
Si I	+0.07	-0.12	+0.05
K I	+0.10	+0.10	+0.10
Ca I	-0.10	+0.01	+0.15
Sc II	0.00	-0.15	+0.10
Ti I	-0.17	-0.01	+0.09
V II	-0.15	-0.03	+0.17
Cr I	-0.12	+0.03	+0.13
Cr II	+0.09	-0.18	+0.05
Mn I	+0.05	+0.05	0.00
Fe I	-0.01	-0.08	+0.13
Fe II	+0.14	-0.24	+0.07
Co I	-0.01	-0.10	+0.11
Ni I	0.00	-0.10	+0.16
Cu I	+0.10	0.00	+0.20
Zn I	+0.12	-0.16	+0.05
Rb I	-0.15	0.00	0.00
Y I	-0.19	0.00	+0.07
Y II	-0.01	-0.13	+0.07
Zr I	-0.10	+0.05	0.00
Ba II	0.00	-0.10	+0.25
La II	-0.02	-0.12	+0.08
Nd II	-0.03	-0.11	+0.02
Eu II	0.00	-0.12	0.00

10%, are available for a large fraction of lines. For other Fe-peak elements, errors in their *gf* values may range between 10 and 25%. For neutron-capture elements the accuracies of recent estimates are in the 10 to 25% range. An extensive list of *gf* values for all important elements can be found in Sumangala Rao, Giridhar, & Lambert (2012).

The sensitivity of the derived abundances to the uncertainties in the model atmosphere parameters




Fig. 2. Observed (filled squares) and synthetic (solid lines) spectra in the $\lambda 5165$ Å C_2 band region for IRAS 16559-2957. The synthetic spectra correspond, from top to bottom, to $\log \varepsilon(C)=7.95$, 7.75 , and 7.55 .

T_{eff} , $\log g$ and ξ_t are summarized in Table 2. We present abundance variations caused by changes in atmospheric parameters of 100 K, 0.25 dex and 0.2 km s^{-1} respectively, relative to the selected model. We have evaluated the total error in the abundance of a given element, by taking the square root of the summation of the squares of the errors associated to T_{eff} , $\log g$ and ξ_t . From the uncertainties listed in Table 2, we find the total absolute uncertainty to range from 0.05 for C to 0.29 for Ba II.

4.2. Spectrum Synthesis Analysis

In this section we discuss the determination of abundances of light elements, CNO. Some lines that are blended or affected by the hyper-fine structure were analyzed by spectral synthesis using the *synth* driver in the MOOG code. We have validated the line list used as follows. We first have used the line list of the corresponding spectral region and fitted the synthesized spectrum with the solar flux atlas of Kurucz et al. (1984). Since many molecular lines are not very strong in the solar spectrum, we repeated the procedure with the spectral atlas of Arcturus (Hinkle et al. 2000) to corroborate the input atomic and molecular data. There are still a few unidentified features that could contribute to the uncertainties.

4.2.1. Carbon

The carbon abundances were derived from the C_2 Swan band lines at $\lambda 4730$ – 4750 Å and $\lambda 5160$ –

5167 Å. An alternate region containing CH molecules located at 4300 – 4310 Å (G-band) has also been used for this purpose. The C_2 lines have been shown to be useful carbon abundance indicators in carbon stars (Zamora et al. 2009) and in Galactic R Coronae Borealis stars (Hema, Pandey, & Lambert 2012). Since IRAS 16559-2957 is an O-rich star ($C/O \sim 0.11$), C being locked up in molecules is important; hence, allowance was made for C in molecules CH, C_2 , CN and CO at the time of synthesizing C_2 and CH. For IRAS 16559-2957 the C abundance has a mean value $\varepsilon(C)=7.75 \pm 0.20$, i.e. the star is carbon deficient.

Figure 2 shows an example of the observed and synthetic spectra in the region around $\lambda 5165$ Å.

4.2.2. Nitrogen

The nitrogen abundances were derived from the CN molecular lines also known as the CN red system because the molecular bands occurs at $\lambda \geq 5800$ Å. In this study we use the CN(5,1) molecular line at $\lambda 6332.18$ Å. The CN(5,1) feature is affected by blends with two lines at $\lambda 6331.95$ Å, one due to Si I and the other due to Fe II. Both features are taken into account in the synthesis. In order to derive the abundance of N is necessary to consider the C abundance previously obtained from the Swan bands.

Figure 3 shows the observed and synthetic spectra of IRAS 16559-2957 in the region $\lambda 6331$ – 6333 Å. We estimate a value of $\log \varepsilon(N)=9.2 \pm 0.3$ from the $\lambda 6332$ Å region. This value is abnormally high and is estimated with larger uncertainty probably due to the fact that the feature is heavily blended and the region contains unidentified lines.

4.2.3. Oxygen

Unfortunately, the [O I] forbidden line at 6300.304 Å region is not present in the observed spectrum of IRAS 16559-2957 as it falls in an inter order gap. The [O I] line at 5577.3 gives a value $\varepsilon(O)=9.0$; however, it is blended with a C_2 feature and we opted for not using it. Thus, we determined the oxygen abundance from the equivalent width of the line 6363.78 Å. However, the derived value of $\varepsilon(O)=8.7$ should be considered an upper limit since this line seems contaminated with either nearby metallic lines or molecular CN.

4.2.4. $^{12}C/^{13}C$

The $^{12}C/^{13}C$ isotope ratio was determined using the features of ^{12}CN and ^{13}CN molecules in the 5622 – 26 Å spectral region (Figure 4). We derived a value of the $^{12}C/^{13}C$ isotope ratio equal to 15.




Fig. 3. Observed (filled squares) and synthetic (solid lines) spectra of the CN(5,1) band. The synthetic spectra correspond, from top to bottom to $\log \epsilon(N) = 9.10, 9.20,$ and 9.30 .

4.2.5. Hyper-fine structure considerations

Hyper-fine structure (hfs) has the effect of desaturating strong lines; hence, it is important to consider hyper-fine components when performing abundance analysis of strong lines (McWilliam et al. 1995). For weak unsaturated lines the hfs treatment is not necessary.

In this analysis we have adopted the hfs data of Spite, Barbuy, & Spite (1989) and Prochaska & McWilliam (2000) for the Sc II lines, Prochaska & McWilliam (2000) and Allen & Porto de Mello (2011) for the Mn I lines, Mucciarelli et al. (2008) for the Eu II line, McWilliam (1998) for the Ba II lines, Allen & Porto de Mello (2011) for the Cu I lines, Reddy et al. (2002) for the Li I doublet at $\lambda 6707.7 \text{ \AA}$ and Lambert & Luck (1976) for the Rb I at 7800.3 \AA .

5. RESULTS AND DISCUSSION

5.1. Derived Elemental Abundances

In Table 3 we present the calculated elemental abundances of IRAS 16559-2957 obtained as discussed in the previous sections⁴. The solar abundances are taken from Asplund, Grevesse, & Sauval (2005).

5.2. Carbon, Nitrogen and Oxygen

In order to infer the evolutionary status of IRAS 16559-2957 the determination of CNO abun-

⁴We adopt the usual notation $[X/H] = \log N(X)/N(H)_* - \log N(X)/N(H)_\odot$, where $\log N(H) \equiv 12$ is the hydrogen abundance by number.




Fig. 4. Determination of the $^{12}\text{C}/^{13}\text{C}$ isotope ratio for IRAS 16559-2957. Observed (filled squares) and synthetic (solid lines) spectra in $\lambda 5622\text{--}28 \text{ \AA}$. Our best estimate is $^{12}\text{C}/^{13}\text{C} = 15$ (green). The colour figure can be viewed online.

dances is required. Table 4 shows the C/O, C/N and $^{12}\text{C}/^{13}\text{C}$ ratios, the abundance of CNO relative to the iron abundance. An average iron abundance of $[\text{Fe}/\text{H}] = +0.04 \pm 0.19$ was obtained from 42 Fe I and 9 Fe II lines.

The peculiarity present in the CN abundances is the deficiency of C ($[\text{C}/\text{Fe}] = -0.68$) and the enhancement of N ($[\text{N}/\text{Fe}] = +1.38$). Our value $\sum \text{CNO}/\text{Fe} = +0.16 \pm 0.2$ is nearly solar, as expected before the third dredge up (Iben 1964) and being the O abundance also solar, the C deficiency and its uncertainty impact on the N abundance. The uncertainty in the C abundance of ± 0.2 indicates that C might be underestimated by a factor of 1.6. This and the non-LTE effect may produce an overestimation of N by as much as 0.3–0.4 dex; however, the N enhancement is still significant.

Strong C deficiency and N enhancement are the consequence of CN-cycled material dredged up to the surface from the He-burning shell (first dredge up) in stars upon ascending to the red giant branch. It has also been observed in core He-burning stars (Tautvaišienė et al. 2001) and in RHB stars (Afsar, Sneden, & For 2012).

In massive post-AGB stars ($M > 4 M_\odot$) N is also enhanced as in the hot bottom of the convective envelope carbon produced by helium burning is converted into nitrogen (Lattanzio & Wood 2004; Giridhar 2011). This keeps the C/O ratio below unity

TABLE 3

ELEMENTAL ABUNDANCES FOR
IRAS 16559-2957

Species	$\log \epsilon_{\odot}$	[X/H]	s.d.	N	[X/Fe]
C(C ₂)	8.39	-0.64		syn	-0.68
N(CN)	7.78	+1.42		syn	+1.38
O I	8.66	+0.04		1	0.00
Li I	1.05	-0.07		syn	-0.11
Na I	6.17	+0.47	± 0.19	3	+0.43
Mg I	7.53	+0.24	± 0.16	2	+0.20
Al I	6.37	+0.10	± 0.14	4	+0.06
Si I	7.51	+0.05	± 0.13	8	+0.01
K I	5.08	-0.28		syn	-0.32
Ca I	6.31	+0.01	± 0.09	6	-0.03
Sc II	3.05	-0.15		syn	-0.19
Ti I	4.90	-0.27	± 0.12	11	-0.31
V I	4.00	+0.14	± 0.15	14	+0.10
Cr I	5.64	-0.13	± 0.08	2	-0.17
Cr II	5.64	-0.06	± 0.13	3	-0.10
Mn I	5.39	-0.37		syn	-0.41
Fe I	7.45	+0.06	± 0.13	42	
Fe II	7.45	+0.03	± 0.16	9	
Co I	4.92	+0.35	± 0.17	4	+0.31
Ni I	6.23	+0.28	± 0.12	13	+0.24
Cu I	4.21	-0.71		syn	-0.75
Zn I	4.60	+0.08		1	+0.04
Rb I	2.60	-0.20		syn	-0.23
Y I	2.21	-0.27		1	-0.31
Y II	2.21	-0.37		1	-0.41
Zr I	2.59	-0.19		syn	-0.23
Ba II	2.17	-0.62		syn	-0.66
La II	1.13	+0.34	± 0.25	2	+0.30
Nd II	1.45	-0.63		1	-0.67
Eu II	0.52	-0.22		syn	-0.26

and the star does not become C-rich (Boothroyd, Sackmann, & Ahern 1993). This Hot Bottom Burning process also features efficient proton capture and hence s-process element enhancements.

IRAS 16559-2957 is an O-rich (C/O \sim 0.11) and is not s-process enriched. The high nitrogen abundance [N/Fe]=+1.38 clearly exceeds the mean observed values in giant ([N/Fe]=+0.35) and dwarf ([N/Fe]=+0.30) stars (Luck 1991; Reddy, Lambert, & Allende 2006). The O abundance, [O/Fe]=0.0, is similar to those observed in giant stars ([O/Fe] \sim +0.08) (Luck 1991; Mishenina et al. 2006; Lambert & Ries 1981; Bensby, Feltzing, & Lundström 2003).

5.3. Other heavy elements

Other relevant abundances are addressed in this section. The Li abundance shows depletion. Theoretical predictions suggests that the stellar Li abundance, if starting from the present interstellar medium abundance $\log \epsilon(\text{Li})=3.3$, would be down to around 1.5 after the first dredge up. Hence G and K giants are expected to have low lithium abundances (Brown et al. 1989). The value found for IRAS 16559-2957, $\log \epsilon(\text{Li})=0.98 \pm 0.13$ is therefore in reasonable agreement with the first dredge-up calculations and suggests that surface Li abundance has been depleted already during the main sequence evolution.

IRAS 16559-2957 shows relative enrichment of sodium ([Na/Fe]=+0.43). A possible source of error as explained by Mishenina et al. (2006) is non-LTE effect. Our Na abundance was calculated from three lines: 5682, 6154 and 6160 Å, with equivalent widths \sim 135 mÅ for which the corrections would be in the +0.10 to +0.15 range; hence the relative enhancement of Na is real and not caused by the neglect of non-LTE treatment.

The abundances of α elements are, within the uncertainties, of solar value as they are in dwarf and giant stars of the thin disk (Reddy et al. 2003, 2006; Luck & Heiter 2007).

The abundances of iron-peak elements (Sc, V, Mn, Fe, Co, Ni) in IRAS 16559-2957 are consistent with the calculated trends by Takeda, Sato, & Murata (2008) and Wang et al. (2011) for field G-giants, Reddy et al. (2003, 2006) for FG-dwarfs and Reddy, Giridhar, & Lambert (2012) for red giant members of open cluster.

The copper underabundance [Cu/Fe]=-0.75 calls our attention. This is comparable to values in metal-poor field stars (Snedden, Gratton, & Crocker 1991). Expected values for [Cu/Fe] in open cluster giants are between -0.11 to -0.23 while for normal giants of the thin disk the value is expected to be $+0.01 \pm 0.13$ (Reddy et al. 2012; Luck & Heiter 2007). Our value defies a simple explanation.

The abundance of rubidium was obtained from spectral synthesis using the Rb I line at 7800 Å. Our synthesis profile takes into account the hyper-fine structure of the ⁸⁵Rb and ⁸⁷Rb isotopes (Lambert & Luck 1976). The fit to this line includes the Si I line at 7799.99 Å for which we took its gf value from Gustafsson et al. (2008) and we checked the atomic data of the spectral region by fitting the Arcturus spectrum (Hinkle et al. 2000). The obtained abundance value [Rb/Fe]=-0.23 \pm 0.19 can be compared with the results of Carney et al. (2005) for

TABLE 4
CNO ABUNDANCES FOR IRAS 16559-2957

C/O	C/N	$^{12}\text{C}/^{13}\text{C}$	[C/Fe]	[N/Fe]	[O/Fe]	[Fe/H]
0.11 ± 0.08	0.04 ± 0.03	15	-0.68 ± 0.27	$+1.38 \pm 0.42$	0.00 ± 0.19	$+0.04 \pm 0.19$

four field giant stars in the direction of the southern warp of the Galactic disk ($\langle[\text{Rb}/\text{Fe}]\rangle = +0.12 \pm 0.03$). We find that the difference is hardly significant. Our value is also consistent with the mean abundance ratio of six giant stars in the old open clusters Be 20 and Be 29 in the outer Galactic disk ($\langle[\text{Rb}/\text{Fe}]\rangle = -0.09 \pm 0.12$) and of three giant stars in the old open cluster M67 ($\langle[\text{Rb}/\text{Fe}]\rangle = -0.27 \pm 0.05$) (Carney et al. 2005).

In the spectrum of IRAS 16559-2957 we have detected lines of the neutron capture elements Y, Zr, Ba, La, Nd, and Eu. The derived abundances of these elements are similar to those observed in field G-giants (Takeda et al. 2008; Wang et al. 2011), in FG-dwarfs (Reddy et al. 2003, 2006) and in red giant members of open cluster (Reddy et al. 2012). However we note that Ba and Nd are significantly diminished relative to those observed in dwarf and giant stars.

5.4. Discussion

The chemical diversity of the central stars of PAGB's is very large. They spread a large range of temperatures and masses below $8 M_{\odot}$. Different families are described in detail in the thematic review by van Winckel (2003) and more recently by Garca-Lario (2006) and Giridhar (2011). There are two known types of O-rich PAGB's; the so called *classical*, optically bright which exhibit low metallicities ([Fe/H] between -1 and -0.3) and display a double peak in their Spectral Energy Distribution (SED), an optical one from the central star and a near-IR one from the circumstellar material. They do not display s-process enhancements and seem to be the successors of AGB stars that do not experience an efficient third dredge-up (van Winckel 2003).

The second group is formed by more massive stars ($M > 4 M_{\odot}$) where the Hot Bottom Burning burns ^{12}C into ^{13}C and ^{14}N and the products are brought up to the surface via the third dredge-up; this can keep the C/O ratio below unity, maintain the $^{12}\text{C}/^{13}\text{C}$ ratio near the CN equilibrium value of 3.5 and dramatically enhance ^{14}N as explained by Lattanzio (2003). The stars in the later group are surrounded by dusty molecular shells responsible of IR emission and CO and OH masers.

The optical counterpart of IRAS 16559-2957 is an O-rich star, and its most prominent spectral properties found in the present paper are: mild iron deficiency $[\text{Fe}/\text{H}] = +0.04 \pm 0.19$, $\text{C}/\text{O} = 0.11$, $^{12}\text{C}/^{13}\text{C} = 15$, and no signs of s-process enhancements. In fact an indicator of the influence of proton flux is the ratio of heavy to light s-process elements or [hs/ls] (Luck & Bond 1991). In IRAS 16559-2957 we calculate $[\text{hs}/\text{ls}] = -0.04$, where we have taken the abundances of Ba, La and Nd as representatives of the heavy s-process elements and Y and Zr a representative of the light ones. Typical value in s-processed enhanced PAGB's is $[\text{hs}/\text{ls}] \sim 10$ (Reddy et al. 2002).

The isotope ratio $^{12}\text{C}/^{13}\text{C}$ is not by itself an indicator of evolution from AGB to PAGB stages and its value changes dramatically between C-rich and O-rich stars. In C-rich stars a lower limit of $^{12}\text{C}/^{13}\text{C} > 20$ has been found by Bakker et al. (1997) using circumstellar molecular CN red lines, and $^{12}\text{C}/^{13}\text{C} > 25$ found by Reddy et al. (2002) using ^{12}CN and ^{13}CN features in the near-IR. Values in the range $30 < ^{12}\text{C}/^{13}\text{C} < 70$ were found in a sample of carbon stars in the Galactic disk by Lambert et al. (1986). In O-rich stars however, the first dredge up reduces the surface $^{12}\text{C}/^{13}\text{C}$ ratio and in low mass stars the inefficient third dredge up maintains the ratio low. More massive stars with an intense Hot Bottom Burning (HBB) also maintain the isotope ratio low. $^{12}\text{C}/^{13}\text{C}$ ratios as low as 3–5 have been found in O-rich planetary nebulae (Rubin et al. 2004). According to Lattanzio & Forestini (1999) low values of $^{12}\text{C}/^{13}\text{C}$ are strong evidence of HBB while high values are indication of ^{12}C dredge up without HBB and intermediate values are harder to interpret.

Thus, although not an indication of evolution in our O-rich star, our value of $^{12}\text{C}/^{13}\text{C} = 15$ is consistent with first dredge up predictions, accompanied of C decrease and N enhancement (Iben 1965). It is also comparable to the mean ratio $^{12}\text{C}/^{13}\text{C} = 16$ in the sample of mildly metal poor giants of Cottrell & Sneden (1986). Giants in older open clusters, with lower turn-off mass, exhibit values between 10 and 20 (Gilroy & Brown 1991; Smiljanic et al. 2009). The clump giant stars in open clusters also shows low values of $^{12}\text{C}/^{13}\text{C}$ ratios suggesting a non-standard

mixing episode occurring sometime along the upper RGB (Tautvaišienė et al. 2010).

Lithium in IRAS 16559-2957 is depleted. Lithium can be produced in AGB stars through the Cameron-Fowler Mechanism (Cameron & Fowler 1971) and large enhancements have been observed in AGB's in the Magellanic Clouds (Smith & Lambert 1990). This, which is the consequence of HBB, seems to work only in intermediate mass stars ($M > 5 M_{\odot}$) (Lattanzio & Forestini 1999).

In IRAS 16559-2957 it was noted that the Na is enriched showing that the surface abundance of this element has suffered changes at the AGB, likely caused by proton capture on ^{22}Ne during the HBB phase as explained by Lattanzio (2003).

6. CONCLUSIONS

We present the results of a detailed atmospheric abundance analysis of the star IRAS 16559-2957, previously classified as a post-AGB. The analysis is based on a high resolution spectrum of the $V \sim 13$ mag object at the coordinates of IRAS 16559-2957 and includes 27 elements. The photospheric chemical abundances are derived by spectral synthesis and from the equivalent widths of selected spectral lines.

Our spectrum is quite different from the low resolution spectrum published by Hu et al. (1993). The derived temperature in the present work, 4250 ± 200 K, corresponds more closely to an early K-type stars and not to an F5 as classified by these authors.

The spectral characteristics of IRAS 16559-2957 do not correspond with the properties of O-rich PAGB's. On the other hand, many of them compare well with properties of giant stars at the AGB as it has been documented in §§§ 5.2, 5.3 and 5.4.

Our results $^{12}\text{C}/^{13}\text{C}=15$ and $[\text{C}/\text{Fe}]=-0.68$ are consistent with first dredge up predictions (Iben 1965). We also note the presence of Rb, only observed in AGB stars, very few field giant stars and in old open clusters.

The small photospheric radial velocity of -2.3 ± 0.5 km s $^{-1}$ ($V_{\text{LSR}} = +7.1$ km s $^{-1}$) is typical of the thin disk population. It is supported by the near solar metallicity measured for this object. The OH maser V_{LSR} of 57–70 km s $^{-1}$ (te Lintel Hekkert et al. 1991) corresponds to the material being ejected, hence red shifted to the observer. Our high resolution spectrum, and the low resolution spectrum of Hu et al. (1993) do not exhibit any emission, hence circumstellar material causing IR flux and maser may be much farther from the photosphere and cooler. OH masers in AGB stars are common (e.g., Reid et al. 1977; Bowers, Johnston, & de Veig

1989) and occur at distances $>10^3$ AU from the central star (Reid & Moran 1981). No OH maser blue shifted component has been observed. If this scenario is correct, given the low observed stellar velocity, either the outflow is not spherically symmetric or the blue component should have a velocity of ~ -60 km s $^{-1}$, which would indicate a very large outflow velocity, but this has been seen in some post-AGB shells, e.g., Zijlstra et al. (2001).

From the atmospheric detailed abundance analysis, we conclude that IRAS 16559-2957 is experiencing its first crossing towards the red giant branch and has probably undergone the first dredge-up. No traces of third dredge up are observed. IRAS 16559-2957 is unlikely to be a post-AGB star.

We are indebted to Prof. D. L. Lambert for permitting the use of the high resolution spectrum of IRAS 16559-2957 used in this work and to Prof. Sunetra Giridhar for her comments and suggestions on the abundance analysis. The pertinent comments and input of an anonymous referee are fully acknowledged. This work was supported by DGAPA-Universidad Nacional Autónoma de México project IN104612.

REFERENCES

- Afsar, M., Sneden, C., & For, B. Q. 2012, *AJ*, 144, 20
 Allen, D. M., & Porto de Mello, G. F. 2011, *A&A*, 525, A63
 Alonso, A., Arribas S., & Martínez-Roger, G. 1999, *A&AS*, 140, 261
 Asplund, M., Grevesse, N., & Sauval, A. J. 2005, in *ASP Conf. Ser. 336, Cosmic Abundances as Records of Stellar Evolution and Nucleosynthesis in Honor of David L. Lambert*, ed. T. G. Barnes III & F. N. Bash (San Francisco: ASP), 25
 Bakker, E. J., van Dishoeck, E. F., Waters, L. B. F. M., & Schoenmaker, T. 1997, *A&A*, 323, 469
 Bensby, T., Feltzing, S., & Lundström, I. 2003, *A&A*, 410, 527
 Bonifacio, P., Caffau, E., & Molaro, P. 2000, *A&AS*, 145, 473
 Boothroyd, A. I., Sackmann, I. J., & Ahern, S. C. 1993, *ApJ*, 416, 762
 Boyarchuck, A. A., Lyubimkov, L. S., & Sakhbullin, N. A. 1985, *Astrophysics*, 22, 203
 Bowers, P. F., Johnston, K. J., & de Veig, C. 1989, *ApJ*, 340, 479
 Brown, J. A., Sneden, C., Lambert, D. L., & Dutchover, E., Jr. 1989, *ApJS*, 71, 293
 Cameron, A. G. W., & Fowler, W. A. 1971, *ApJ*, 164, 111
 Carney, B. W., Yong, D., Teixeira de Almeda, M. L., & Seitzer, P. 2005, *AJ*, 130, 1111
 Castelli, F., & Kurucz, R. L. 2003, in *IAU Symp. 210, Modelling of Stellar Atmosphere*, ed. N. E. Piskunov, W. W. Weiss & D. F. Gray (San Francisco: ASP), 20
 Cottrell, P. L., & Sneden, C. 1986, *A&A*, 161, 314

- Cutri, R. M., et al. 2003, The IRSA 2MASS All-Sky Point Source Catalog of Point Sources, NASA/IPAC Infrared Science Archive
- García-Lario, P. 2006, in IAU Symp. 234, Planetary Nebulae in our Galaxy and Beyond, ed. M. J. Barlow & R. H. Méndez (Cambridge: Cambridge Univ. Press), 63
- García-Lario, P., Manchado, A., Pych, W., & Pottasch, S. R. 1997, *A&AS*, 126, 479
- Gilroy, K. K., & Brown, J. A. 1991, *ApJ*, 371, 578
- Giridhar, S. 2011, in ASI Conference Ser. 3, ed. P. Khare & C. H. Ishwara-Chandra, 39
- Gustafsson, B., Edvardsson, B., Eriksson, K., Jørgensen, U. G., Nordlund, A., & Plez, B. 2008, *A&A*, 486, 951
- Hema, B. P., Pandey, G., & Lambert, D. L. 2012, *ApJ*, 747, 102
- Hinkle, K., Wallace, L., Valenti, J., & Harmer, D. 2000, Visible and Near Infrared Atlas of the Arcturus Spectrum 3727–9300 Å, ed. K. Hinkle, L. Wallace, J. Valenti, & D. Harmer (San Francisco: ASP)
- Hu, J. Y., Slijkhuis, S., de Jong, T., & Jiang, B. W. 1993, *A&AS*, 100, 413
- Kurucz, R. L., Furenlid, I., Brault, J., & Testerman, L. 1984, Solar Flux Atlas from 296 to 1300 nm (New Mexico: National Solar Observatory)
- Iben, I. 1964, *ApJ*, 140, 1631
- _____. 1965, *ApJ*, 142, 1447
- Lambert, D. L., Gustafsson, B., Eriksson, K., & Hinkle, K. H. 1986, *ApJS*, 62, 373
- Lambert, D. L., & Luck, R. E. 1976, *Obs*, 96, 100
- Lambert, D. L., & Ries L. M. 1981, *ApJ*, 248, 228
- Lattanzio, J. 2003, in IAU Symp. 209, Planetary Nebulae: Their Evolution and Role in the Universe, ed. S. Kwok, M. Dopita, & R. Sutherland (San Francisco: ASP), 73
- Lattanzio, J., & Forestini, M. 1999, in IAU Symp. 191, Asymptotic Giant Branch Stars, ed. T. Le Bertre, A. Lébre, & C. Wealkens (San Francisco: ASP), 31
- Lattanzio, J., & Wood, P. R. 2004, in Asymptotic Giant Branch Stars, ed. H. J. Habing & H. Olofsson (New York: Springer-Verlag), 22
- Luck, R. E. 1991, *ApJS*, 75, 579
- Luck, R. E., & Bond, H. E. 1991, *ApJS*, 77, 515
- Luck, R. E., & Heiter, U. 2007, *AJ*, 133, 2464
- McWilliam, A. 1998, *AJ*, 115, 1640
- McWilliam, A., Preston, G. W., Sneden, C., & Searle, L. 1995, *AJ*, 109, 2757
- Mishenina, T. V., Bienaymé, O., Gorbaneva, T. I., Charbonnel, C., Soubiran, C., Korotin, S. A., & Kovtyukh V. V. 2006, *A&A*, 456, 1109
- Mucciarelli, A., Caffau, E., Freytag, B., Ludwing, H. G., & Bonifacio, P. 2008, *A&A*, 484, 841
- Prochaska, J. X., & McWilliam, A. 2000, *AJ*, 537, 57
- Ramos-Larios, G., Guerrero, M. A., Suárez, O., Miranda, L. F., & Gómez, J. F. 2009, *A&A*, 501, 1207
- Reddy, A. B. E., Lambert, D. L., & Allende Prieto, A. 2006, *MNRAS*, 367, 1329
- Reddy, A. B. E., Lambert, D. L., Laws, C., González, G., & Covey, K. 2002, *MNRAS*, 335, 1005
- Reddy, A. B. E., Tomkin, J., Lambert, D. L., & Allende Prieto, C. 2003, *MNRAS*, 340, 304
- Reddy, A. B. S., Giridhar, S., & Lambert, D. L. 2012, *MNRAS*, 419, 1350
- Reid, M. J., & Moran, J. M. 1981, *ARA&A*, 19, 231
- Reid, M. J., Muhleman, D. O., Moran, J. M., Johnston, K. J., & Schwartz, P. R. 1977, *ApJ*, 214, 60
- Rubin, R. H., Ferland, G. J., Chollet, E. E., & Horstmeier, R. 2004, *ApJ*, 605, 784
- Sahin, T., & Lambert, D. L. 2009, *MNRAS*, 398, 1730
- Schiller, F., & Przybilla, N. 2008, *A&A*, 479, 849
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Smiljanic, R., Gauderon, R., North, P., Barbuy, B., Charbonnel, C., & Mowlavi, N. 2009, *A&A*, 502, 267
- Smith, V. V., & Lambert, D. L. 1990, *ApJ*, 361, L69
- Snedden, C. 1973, PhD Thesis, Univ. Texas, USA
- Snedden, C., Gratton, R. G., & Crocker, D. A. 1991, *A&A*, 246, 354
- Spite, M., Barbuy, B., & Spite, F. 1989, *A&A*, 222, 35
- Suárez, O., García-Lario, P., Manchado, A., Manteiga, M., Ulla, A., & Pottasch, S. R. 2006, *A&A*, 458, 173
- Sumangala Rao, S., Giridhar, S., & Lambert, D. L. 2012, *MNRAS*, 419, 1254
- Szczerba, R., Siódmiak, N., Stasińska, G., & Borkowski, J. 2007, *A&A*, 469, 799
- Szczerba, R., et al. 2012, in IAU Symp. 283, Planetary Nebulae: An Eye to the Future, ed. A. Manchado, L. Stanghellini, & D. Schönberner (Cambridge: Cambridge Univ. Press), 506
- Takeda, Y., Sato, B., & Murata, D. 2008, *PASJ*, 60, 781
- Tautvaišienė, G., Edvardsson, B., Puzeras, E., Barisevičius, G., & Ilyin I. 2010, *MNRAS*, 409, 1213
- Tautvaišienė, G., Edvardsson, B., Tuominen, I., & Ilyin, I. 2001, *A&A*, 380, 578
- te Lintel Hekkert, P., Caswell, J. L., Habing, H. J., Haynes, R. F., & Norris, R. P. 1991, *A&AS*, 90, 327
- Thévenin, F., & Idiart, T. P. 1999, *ApJ*, 521, 753
- Tull, R. G., MacQueen, P. J., Sneden, C., & Lambert, D. L. 1995, *PASP*, 107, 251
- van der Veen, W. E. C. J., & Habing, H. J. 1988, *A&A*, 194, 125
- van Winckel, H. 2003, *ARA&A*, 41, 391
- Wang, L., Liu, Y., Zhao, G., & Sato, B. 2011, *PASJ*, 63, 1035
- Zamora, O., Abia, C., Plez, B., Domínguez, I., & Cristallo, S. 2009, *A&A*, 508, 909
- Zijlstra, A. A., te Lintel Hekkert, P., Pottasch, S. R., Caswell, J. R., Ratag, M., & Habing, H. J. 1989, *A&A*, 217, 157
- Zijlstra, A. A., et al. 2001, *MNRAS*, 322, 280

R. E. Molina: Laboratorio de Investigación en Física Aplicada y Computacional, Universidad Nacional Experimental del Táchira, Venezuela, (rmolina@unet.edu.ve).

A. Arellano Ferro: Instituto de Astronomía, Universidad Nacional Autónoma de México, Apdo. Postal 70-264, 04510, México, D.F., Mexico (armando@astro.unam.mx).