

A CROSS-CHECK FOR H_0 FROM LYMAN- α FOREST AND BARYON ACOUSTIC OSCILLATIONS

V. C. Busti,^{1,3} R. N. Guimarães,^{2,4} and J. A. S. Lima³

Received September 1 2014; accepted October 13 2015

RESUMEN

Se propone un nuevo método para inferir el valor de la constante de Hubble, H_0 , mediante el flujo medio observado de cuásares de alto corrimiento al rojo y las oscilaciones acústicas de los bariones (BAOs). Se adopta un modelo semianalítico para la distribución de densidad volumétrica independiente de la cosmología que nos permite obtener límites para los parámetros cosmológicos una vez que se cuenta con una aceptable estimación de los parámetros del medio intergaláctico (IGM). Basado en dos muestras diferentes del bosque de Lyman- α y en la medición de las BAOs, nuestro análisis restringe (h, Ω_m) a los intervalos $0.19 \leq \Omega_m \leq 0.23$ y $0.53 \leq h \leq 0.82$. Si bien estos límites son menos estrictos que los que se obtienen por otros métodos, señalamos que con muestras más numerosas y con un mejor conocimiento del IGM nuestro método podría aportar resultados complementarios para la determinación de la constante de Hubble, independientemente de la escala cósmica de distancias.

ABSTRACT

A new method is proposed to infer the Hubble constant H_0 through the observed mean transmitted flux from high-redshift quasars and the baryon acoustic oscillations (BAOs). A semi-analytical model for the cosmological-independent volume density distribution function was adopted; it allowed us to obtain constraints on the cosmological parameters once a moderate knowledge of the Inter Galactic Medium (IGM) parameters is assumed. Our analysis, based on two different samples of Lyman- α forest and the BAO measurement, restricts (h, Ω_m) to the intervals $0.19 \leq \Omega_m \leq 0.23$ and $0.53 \leq h \leq 0.82$ (1σ). Although the constraints are weaker compared with other estimates, we point out that, with a bigger sample and a better knowledge of the IGM, this method could provide complementary results to measure the Hubble constant independently of the cosmic distance ladder.

Key Words: cosmological parameters — distance scale — intergalactic medium

1. INTRODUCTION

Measurements of the Hubble constant H_0 are crucial for the establishment of a cosmic concordance model in cosmology. This constant plays a role in most of cosmic calculations, such as those about the age of the Universe, its size and energy density, pri-

mordial nucleosynthesis, the physical distances between objects, etc. Its importance will even increase in the next decade, since new missions and observational projects are currently being designed to provide accurate measurements of the Hubble constant, e.g., improved parallax calibration for Galactic Cepheids with *Spitzer* and GAIA, and 1% precision of H_0 with *LISA* based on the gravitational radiation from inspiraling massive black holes, which would also increase the precision of the other cosmological parameters.

There are several ways to determine H_0 by using low and high-redshift sources. The most common methods are based on the period-luminosity re-

¹Astrophysics, Cosmology & Gravity Centre (ACGC), and Department of Mathematics & Applied Mathematics, University of Cape Town, South Africa.

²CAPES Foundation, Ministry of Education of Brazil, BrasíliaDF, Brazil.

³Instituto de Astronomia, Geofísica e Ciências Atmosféricas (IAG-USP), Universidade de São Paulo, São Paulo, Brazil.

⁴Biomedical Informatics Group (IBIME), ITACA Institute, Technical University of Valencia, Valencia, Spain.

lation of cepheids, the tip of the red giant branch (TRGB) and the type Ia supernovae (SNe Ia) observed in the local Universe (Freedman & Madore 2010). However, some alternative procedures, such as the Sunyaev–Zel’dovich effect combined with X-ray emission from clusters, which allow a measurement of H_0 from high-redshift objects, have also been discussed (Cunha et al. 2007; Holanda et al. 2012).

Actually, independent estimates are needed in order to obtain a reliable value for H_0 that is not plagued by systematic errors arising from astrophysical environments, as well as from calibrations related to the cosmic distance ladder. In this regard, the importance and cosmological interest of different estimates of H_0 that are independent of the distance ladder has also been discussed by many authors (see Jackson 2007, and references therein for a recent review).

Recently, a tension was reported between the value obtained by (Riess et al. 2011) of $H_0 = 73.8 \pm 2.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$ (1σ), based on measurements of cepheids and SNe Ia (or alternatively, that of Freedman et al. 2012, who obtained $H_0 = 74.3 \pm 2.1 \text{ km s}^{-1} \text{ Mpc}^{-1}$ (1σ), and the value released by *Planck* of $H_0 = 67.3 \pm 1.2 \text{ km s}^{-1} \text{ Mpc}^{-1}$ (1σ), Ade et al. 2014), obtained from measurements of the anisotropies of the cosmic microwave background (CMB). The former is based on local measurements, while the latter assumes a cosmological model to infer H_0 . There are several possible explanations of this discrepancy, such as the following: we live in a Hubble bubble and the local value of H_0 is subject to cosmic variance (Marra et al. 2013); or there is a problem in the calibration of the cepheids and the true value of H_0 is lower, as given by (Sandage et al. 2006); or with the TRGB and SNe Ia given by (Tammann & Reindl 2013); or an extension beyond the standard model of cosmology is needed.

In fact, it is the last option which must be considered, since the emergence of the Λ CDM model as the concordance model in 1998, based on the SNe Ia observations (Riess et al. 1998; Perlmutter et al. 1999), brought with it the cosmological constant and the coincidence problems. As a complete and convincing solution to these problems is still unknown, it is very important to test all the hypotheses of the model. In addition, it is important to verify that different analyses give the same results, since several kinds of data covering different cosmic epochs are available.

In this regard, a possible alternative procedure to measure H_0 is to consider the Lyman- α ($\text{Ly}\alpha$) forest data and the acoustic peak caused by the baryons acoustic oscillations. The $\text{Ly}\alpha$ forest probes the low density intergalactic medium (IGM) over a unique range of redshifts and environments, as seen in the spectra of high-redshift quasars. The observed mean transmitted flux depends on the local optical depth, which in turn depends on the expansion rate of the Universe $H(z)$ and some properties characterizing the intervening medium along our line of sight to any quasar. In principle, the values of some cosmological parameters, including the Hubble constant, could be constrained if we had a moderate knowledge of the IGM properties, such as the hydrogen photoionization rate and the mean temperature, and combined this information with the BAO measurement.

In the last decade, several studies have been performed using $\text{Ly}\alpha$ forests as cosmological tools (McDonald et al. 2000; Weinberg et al. 2003; Wyithe et al. 2008; Viel et al. 2009). Some works use the flux power spectrum of the $\text{Ly}\alpha$ forest to infer the cosmological parameters in two ways: inverting the flux power spectrum to estimate the underlying dark matter power spectrum (Croft et al. 1998, 1999; Hui 1999; Nusser & Haehnelt 1999), or using the power spectrum directly (McDonald et al. 2000; Mandelbaum et al. 2003; McDonald et al. 2005). More recently, new possibilities for using $\text{Ly}\alpha$ forests for cosmological purposes were created by the detection of baryon acoustic oscillations in the correlation function of the transmitted flux fraction (Busca et al. 2013; Slosar et al. 2013).

A simpler approach is to use a semi-analytical model to describe the IGM as proposed by (Miralda-Escudé et al. 2000), who derived a fitting formula for the distribution function of volume density in agreement with simulations that yielded similar results for different numerical methods and different cosmologies (Rauch et al. 1997). This allowed to estimate a theoretical mean transmitted flux and to compare it with observational data in order to constrain the cosmological parameters.

Assuming a flat Λ CDM cosmology, we applied the above described procedure for deriving cross-check values for the Hubble constant H_0 and the matter density parameter Ω_m . Based on two independent samples of $\text{Ly}\alpha$ forest data compiled by (Bergeron et al. 2004) and (Guimarães et al. 2007), respectively, we performed two distinct statistical analyses. First, we considered only the $\text{Ly}\alpha$ forest data set, while the second approach involved a joint analysis combining the $\text{Ly}\alpha$ forest data with the

measurements of the baryon acoustic peak from the WiggleZ survey (Blake et al. 2012). As we shall see, the degeneracy in the free parameters defining the (h, Ω_m) plane is naturally broken when the Ly α data are combined with standard ruler datasets as given by the baryon acoustic oscillations, thereby providing a new independent method to estimate the Hubble constant.

2. THE FOREST SAMPLES

As remarked earlier, in the present analysis we considered two different samples. The first sample consisted of 18 high-resolution high signal-to-noise ratio (S/N) spectra ($2.2 \leq z_{em} \leq 3.3$). The spectra were obtained with the ultraviolet and Visible Echelle Spectrograph (UVES) mounted on the ESO KUEYEN 8.2 m telescope at the Paranal observatory for the ESO-VLT Large Programme (LP) 'Cosmological evolution of the Inter Galactic Medium' (Bergeron et al. 2004). The spectra were taken from the European Southern Observatory archive and are publicly available. This program was devised for gathering a homogeneous sample of echelle spectra of 18 QSOs, with uniform spectral coverage, resolution and signal-to-noise ratio suitable for studying the Ly α forest in the redshift range 1.8 - 3.1. The spectra had a signal-to-noise ratio of 40 to 80 per pixel and a spectral resolution of $\lambda/\Delta\lambda \sim 45000$ in the Ly α forest region. The details of the procedure used for data reduction can be found in (Chand et al. 2004) and (Aracil et al. 2004). All possible metal lines and Lyman systems were flagged and removed from the Lyman forest in this sample. We confirmed that the mean transmitted flux values obtained were consistent with previous works that have used the same data-set (Chand et al. 2004; Aracil et al. 2004; Rollinde et al. 2013).

The second sample consisted of 45 quasars in the redshift range $3.8 \leq z_{em} \leq 4.6$ (Guimarães et al. 2007), where the spectra were measured with the ESI (Sheinis et al. 2002) mounted on the Keck II 10-m telescope. The spectral resolution was $R \sim 4300$, with a signal-to-noise ratio usually larger than 15 per 10 km s $^{-1}$ pixel. This sample has also been studied by Prochaska and colleagues (Prochaska et al. 2003a,b) in their discussions related to the existence of damped Ly α systems. In the present analysis, we used the Ly α forest over the rest-wavelength range 1070-1170 Å, corresponding to an interval of absorption redshift of $3.3 \leq z_{em} \leq 4.4$, in order to avoid contamination by the proximity effect close to the Ly α emission line and possible absorbers associated with OVI. We also carefully avoided regions flagged because of damped/sub-damped Ly α

absorption lines. We used only spectra with mean SNR ≥ 25 . We rejected the broad absorption line QSOs (BAL) and QSOs with more than one damped Ly α system (DLA) redshifted between the Ly- α and the Ly- β emission lines, as their presence may over-pollute the Ly α forest. Metal absorptions were not subtracted from the spectra. We estimate that the number of intervening CIV and MgII systems with $W_{obs} > 0.25 \text{ \AA}$ is of the order of five along each of the lines of sight (e.g. Aracil et al. 2004; Tytler et al. 2004). This means that we expect an error in the determination of the mean absorption of the order of 1 %. This is at least five times smaller than the error expected from the placement of the continuum. Normalization of the spectra is known to be a crucial step in these studies. An automatic procedure (Aracil et al. 2004) iteratively estimates the continuum by minimizing the sum of a regularization term (the effect of which is to smooth the continuum) and a χ^2 term. In our case, the regularization matrix is simply the unit matrix. The procedure smooths the continuum to a fixed scale and the χ^2 term is computed from the difference between the quasar spectrum and the continuum estimated during the previous iteration. The effect of the procedure is that the errors increase when the difference between the spectrum and the continuum is large (e.g., when absorption occurs), so that the weights of the corresponding pixels in the χ^2 minimization is reduced. A smoothing parameter allows to vary the scale of the estimates. This procedure is very successful in removing the strong absorption lines from a local estimate of the continuum and works very well at low redshift. At high redshift, however, the method underestimates the overall absorption in the Lyman- α forest.

We estimated a median error of 5% in the continuum in the whole Ly α forest region at high-redshift. In some regions, the placement can be more uncertain due to the blend of strong lines. However, close to the emission redshift of the QSO, and due to the proximity effect, the Ly α forest is not that dense and the continuum is better defined. The pixels affected by the largest errors are (mostly) those with larger depths (blends of strong lines). Therefore they do not strongly influence the estimation of the transmitted flux.

The mean transmitted flux for the two samples was calculated as the mean flux of all pixels in a bin. At each redshift, we then averaged the transmitted fluxes over all spectra covering this redshift. Figure 1 shows the mean transmitted flux for the two samples. Notice that the errors were calculated us-




Fig. 1. Mean transmitted flux as a function of redshift for $\Omega_m = 0.23$, $\Gamma = 0.8 \times 10^{-12} \text{ s}^{-1}$, $T_0 = 2.5 \times 10^4 \text{ K}$ and some selected values of the h parameter. The data points correspond to the Ly α forest measurements obtained by (Rollinde et al. 2013) (red circles) and (Guimarães et al. 2007) (black squares). The color figure can be viewed online.

ing a bootstrap estimator, even though we only give the diagonal terms in the covariance matrix. The large scatter in the results can be explained by errors in the placement of the continuum fluctuations in the forest number density, cosmic variance, redshift determination, etc. The evolution of the mean transmission with z in the two samples is consistent with previous determinations (e.g. Rauch et al. 1997; Aracil et al. 2004; Tytler et al. 2004; Kirkman et al. 2005; Faucher-Giguère et al. 2008b; Aghaee et al. 2010).

3. BASIC EQUATIONS

The basic measurement obtained from the Ly α forest is the mean transmitted flux at redshift z defined as

$$\langle F \rangle(z) \equiv \langle \exp(-\tau(z)) \rangle, \quad (1)$$

where $\tau(z)$ is the local optical depth, and angle brackets denote an average over the line of sight. Following standard methods (Hui & Gnedin 1997), this paper assumes photoionization equilibrium and a power law temperature-density relation, $T = T_0(1 + \delta)^\beta$, for the low-density IGM, where δ stands for the local overdensity and T_0 for the IGM temperature at mean density ($\delta = 0$). The local optical depth is given by (Peebles 1993; Padmanabhan 2002; Faucher-Giguère et al. 2008a)

$$\tau(z) = A(z)(1 + \delta)^{2-0.75\beta}, \quad (2)$$

with

$$A(z) \equiv \frac{\pi e^2 f_{Ly\alpha}}{m_e \nu_{Ly\alpha}} \left(\frac{\rho_{\text{crit}} \Omega_b}{m_p} \right)^2 \frac{X(X + 0.5Y)}{H(z)} \times \\ \times \frac{R_0 T_0^{-0.75}}{\Gamma} (1 + z)^6.$$

In the expression above, $f_{Ly\alpha}$ is the oscillator strength of the Ly α transition, $\nu_{Ly\alpha}$ is its frequency, m_e and m_p are the electron and proton masses respectively, Ω_b is the baryon density parameter, $H(z)$ is the Hubble parameter, ρ_{crit} is the critical density, X and Y are the mass fractions of hydrogen and helium respectively, which are taken to be 0.75 and 0.25 (Burles et al. 2001), $R_0 = 4.2 \times 10^{-13} \text{ cm}^3 \text{ s}^{-1} / (10^4 \text{ K})^{-0.75}$, and Γ is the hydrogen photoionization rate. The effect of redshift-distortions due to thermal broadening and peculiar velocities on the above equations was neglected since the mean optical depth is in first order independent on these distortions. The mean transmitted flux is a non-linear function of $\langle \tau \rangle$, but the distortions are responsible for only a small effect (Faucher-Giguère et al. 2008b).

The mean transmitted flux is obtained by integrating the local optical depth through a volume density distribution function for the gas $\Delta \equiv 1 + \delta$,

$$\langle F \rangle(z) = \int_0^\infty d\Delta P(\Delta; z) \exp[-\tau(z)]. \quad (3)$$

As it is widely known, Miralda-Escudé et al. (2000) derived an approximate analytical functional form for the distribution function given by

$$P(\Delta; z) = A \exp \left[-\frac{(\Delta^{-2/3} - C_0)^2}{2(2\delta_0/3)^2} \right] \Delta^{-b}, \quad (4)$$

where the parameters A and C_0 are derived by requiring the total volume and mass to be normalized to unity. They extrapolated the distribution function and obtained $\delta_0 = 7.61/(1+z)$ with an accuracy better than 1%, from fits to a numerical simulation at $z = 2, 3$, and 4 from Miralda-Escudé et al. (1996). In order to apply this formalism to our data, the values of b were derived from a cubic interpolation of the values in the simulation. The application of this distribution function to constrain cosmological parameters is well justified, since simulations using different numerical methods and different cosmologies yielded very similar results (Rauch et al. 1997), even though the two simulations share approximately the same amplitude of density fluctuations at the Jeans

scale. However, one must be cautious about the limitations of this distribution function. Actually, this function may be affected by several effects. To cite a few: the numerical properties of the simulation, the thermal state of the gas, fluctuations in the UV background, inhomogeneous reionization of hydrogen and helium (Faucher-Giguère et al. 2008b, and references therein).

From now on we will assume a flat Λ CDM cosmology. In this case, the Hubble parameter is given by

$$H(z) = H_0 [\Omega_m(1+z)^3 + (1 - \Omega_m)]^{1/2}, \quad (5)$$

where we will adopt the convention $h = H_0/100 \text{ km s}^{-1} \text{ Mpc}^{-1}$, which is the Hubble constant normalized in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

Figure 1 shows the effects of H_0 on the mean transmitted flux, along with our observational data (red circles) and the data from (Guimarães et al. 2007) (black squares). From our sample we obtained a scattered data set, compared to the other Ly α sample, which is related to the sample size. It is also evident that our data cover most values of the h parameter; thus, we do not expect tight constraints from the Ly α data set alone.

4. ANALYSIS AND RESULTS

Let us now perform a statistical analysis to find the constraints on the cosmological parameters. The full set of parameters are represented by $\mathbf{p} \equiv (h, \Omega_m, \Gamma, T_0, \Omega_b h^2, \beta)$. We fixed the value of $\Omega_b h^2 = 0.0218$ using the latest observations of deuterium (Pettini et al. 2008) from the Big Bang Nucleosynthesis (Simha & Steigman 2008). We choose $\beta = 0.3$ due to the photoheating during HeII reionization (McQuinn et al. 2009), which is expected to occur in the redshifts covered by our samples. An early reionization model with $\beta = 0.62$ (Hui & Gnedin 1997) yielded similar results. The posterior probability of the parameters $P(\mathbf{p}|d)$ given the data d is

$$P(\mathbf{p}|d) = \frac{P(d|\mathbf{p})P(\mathbf{p})}{P(d)}, \quad (6)$$

where $P(d)$ is a normalization constant, $P(\mathbf{p})$ is the prior over the parameters and $P(d|\mathbf{p}) \propto e^{-\chi^2/2}$ is the likelihood, using the usual definition. For the Ly α forest data,

$$\chi_{Ly\alpha}^2 = \sum_i \frac{(\langle F_{th} \rangle(z_i; \mathbf{p}) - \langle F \rangle(z_i))^2}{\sigma_{\langle F \rangle(z_i)}^2}, \quad (7)$$

where $\langle F_{th} \rangle(z_i; \mathbf{p})$ is the theoretical mean transmitted flux, $\langle F \rangle(z_i)$ is the observational mean transmitted flux and $\sigma_{\langle F \rangle(z_i)}$ is their respective uncertainty. We treat the combination $\Gamma T_0^{0.75}$ as a nuisance parameter with a flat prior. The ranges chosen are $T_0 = [2, 2.5] \times 10^4 \text{ K}$, from estimates of (Zaldarriaga et al. 2001), and $\Gamma = [0.8, 1] \times 10^{-12} \text{ s}^{-1}$ which cover some measurements reported in the literature (Rauch et al. 1997; McDonald & Miralda-Escudé 2001; Meiksin & White 2004; Tytler et al. 2004; Bolton et al. 2005; Kirkman et al. 2005), although they are in disagreement with the values obtained by (Faucher-Giguère et al. 2008a,b), which used a redshift-dependent relation for T_0 not favoured by our data. We defined the 1, 2 and 3 σ confidence intervals as the iso- χ^2 regions given by $\Delta\chi^2 = \chi^2 - \chi_{min}^2$ equals to 2.3, 6.17 and 11.8, respectively.

It is worth noting that there seems to be a tension between the two Ly α samples according to Fig. 1, which shows that the low redshift sample prefers lower values of h . Therefore, we included two parameters to multiply the errors, one for each sample, in order to achieve an acceptable fit. We considered flat priors within the range [1.0, 1.1] and marginalized over them.

In what follows, we first consider the Ly α forest data separately, and, then, we present a joint analysis including the BAO signature extracted from the WiggleZ survey (Blake et al. 2012).

4.1. Limits from the Ly α forest data set

Figure 2 shows the results of the statistical analysis performed with the Ly α forest data. The data do not provide good constraints on both parameters. Several sanity checks were performed in order to validate the values obtained in the statistical analysis. The reduced χ^2 obtained is 0.96. Different values for β , different intervals for $\Gamma T_0^{0.75}$ in the marginalization, or even a redshift-dependence for T_0 , have resulted in poorer fits compared to what is shown in Figure 2. In addition, even when considering that cosmological constraints are affected by the physical parameters of the IGM, the opposite is also true, that is, that different cosmologies may provide results not compatible with the IGM parameters. In this context, it is recommended to fit cosmological and IGM parameters together in order to obtain more reliable results. The interval for $\Gamma T_0^{0.75}$, selected in the marginalization, may suffer from a circularity problem, i.e., the values obtained for T_0 and Γ were obtained assuming a particular cosmological model. Nonetheless, such parameters should reflect local physics, which are not expected to depend




Fig. 2. Confidence regions (68.3%, 95.4% and 99.7%) in the (Ω_m, h) plane provided by Ly α forest data. The basic conclusion is that the Ly α forest alone cannot constrain both parameters.

strongly on the assumed model. With this in mind, the effect of a broader interval on the marginalization is a broader interval in the (Ω_m, h) plane, but with a well-defined slope. Independent estimates of $\Gamma T_0^{0.75}$ are of great interest; Γ may be determined from the proximity effect (Guimarães et al. 2007; Carswell et al. 1982; Murdoch et al. 1986; Rollinde et al. 2005) and T_0 from linewidth measurements, providing an important cross check for the method developed in this work.

Despite the considerations discussed above, Figure 2 shows that the slope in the (h, Ω_m) plane suggests that a joint analysis with an independent test constraining only Ω_m could provide interesting limits to the Hubble constant. Thus, a joint analysis with the BAO data is presented in the next subsection.

4.2. The Ly α forest and BAO: A Joint Analysis

In order to obtain better constraints on the cosmic parameters, we performed a joint analysis involving Ly α forest data and data from baryon acoustic oscillations obtained from the WiggleZ survey (Blake et al. 2012). The BAO scale can be represented by the parameter (Eisenstein 2005)

$$\mathcal{A} \equiv \frac{\Omega_m^{1/2}}{\mathcal{H}(z_*)^{1/3}} \left[\frac{1}{z_*} \Gamma(z_*) \right]^{2/3}, \quad (8)$$

where z_* is the redshift at which the acoustic scale has been measured, $\mathcal{H}(z)$ is $H(z)/H_0$ and $\Gamma(z_*)$ is the dimensionless comoving distance to z_* .




Fig. 3. Contours in the $\Omega_m - h$ plane provided by the joint analysis combining Ly α Forest and BAO. As before, the contours correspond to 68.3%, 95.4% and 99.7% confidence levels. Note that the best-fit model converges to $h = 0.66$ and $\Omega_m = 0.21$. These results should be compared with the ones presented in Figure 2.

From equation (8), it is seen that the BAO scale is independent of h , and thereby yielded constraints only for the matter density parameter. The statistical analysis was performed with $\chi^2 = \chi_{Ly\alpha}^2 + \chi_{BAO}^2$, where

$$\chi_{BAO}^2 = (\mathcal{A}_i - \mathcal{A}_{obs})^T C^{-1} (\mathcal{A}_i - \mathcal{A}_{obs}), \quad (9)$$

\mathcal{A}_i is a vector of three theoretical values at three effective redshifts, \mathcal{A}_{obs} is a vector with the respective observed values and C is the covariance matrix for the observations. Here, we used the data provided in Table 2 by (Blake et al. 2012), where we marginalized over all other parameters.

Figure 3 shows the constraints on the pair of parameters (h, Ω_m) obtained from our joint analysis involving Ly α forest data and BAO. Within a 68.3% confidence level, we obtained $0.19 \leq \Omega_m \leq 0.23$ and $0.53 \leq h \leq 0.82$. The reduced χ^2 is now equals to 1.03. The value for Ω_m is lower than what was found by other studies (Amanullah et al. 2010), but is still consistent within the 68.3% confidence level.

Table 1 shows some recent measurements of H_0 using different techniques and data; we can see that our H_0 value cannot provide stringent limits that would allow us to make a decision about the correct value of H_0 . The interesting aspect is that the method discussed here provides an additional estimate, independent of the cosmic distance ladder and, covering a redshift range ($2 < z < 5$), that other

TABLE 1

LIMITS TO H_0 FROM SEVERAL METHODS. RAND. STANDS FOR RANDOM ERRORS WHILE SYST. FOR SYSTEMATIC ERRORS

Method	Reference	h
Cepheid Variables	(Freedman 2001) (HST Project)	0.72 ± 0.08
Age Redshift	(Jimenez et al. 2003) (SDSS)	0.69 ± 0.12
Age Redshift	(Busti et al. 2014)	0.649 ± 0.042
SNe Ia/Cepheid	(Sandage et al. 2006)	$0.62 \pm 0.013(\text{rand.}) \pm 0.05(\text{syst.})$
SZE+BAO	(Cunha et al. 2007)	$0.74^{+0.04}_{-0.03}$
Old Galaxies + BAO	(Lima et al. 2009)	0.71 ± 0.04
SNe Ia/Cepheid	(Riess et al. 2009)	0.742 ± 0.036
Time-delay lenses	(Paraficz & Hjorth 2010)	0.76 ± 0.03
CMB	(Komatsu et al. 2011) (WMAP7)	0.710 ± 0.025
SNe Ia/Cepheid	(Riess et al. 2011)	0.738 ± 0.024
Median Statistics	(Chen & Ratra 2011)	0.680 ± 0.028
SZE+BAO	(Holanda et al. 2012)	$0.74^{+0.05}_{-0.04}$
CMB	(Hinshaw et al. 2013) (WMAP9)	0.700 ± 0.022
SNe Ia/Cepheid	(Freedman et al. 2012)	0.743 ± 0.021
SNe Ia/TRGB	(Tammann & Reindl 2013)	$0.64 \pm 0.016(\text{rand.}) \pm 0.02(\text{syst.})$
CMB	(Ade et al. 2014) (Planck)	0.673 ± 0.012

techniques cannot provide. Therefore, it is a complementary measurement that can be used as a cross-check with local and global measurements.

5. CONCLUSIONS

In this work, we used a cosmological independent semi-analytical model to describe the IGM and the data from the Ly α forest and baryon acoustic oscillations, in order to constrain cosmological parameters. We provided a cross-check for the Hubble constant H_0 and the matter density parameter Ω_m , which were derived by assuming a flat Λ CDM model; we also established that local properties such as Γ and T_0 are only weakly dependent on the adopted cosmology.

We did not obtain good constrains on both parameters from the statistical analysis using only the Ly α forest data. This happened also due to our scarce knowledge about the local properties of the intergalactic medium. However, when we performed a joint analysis involving the Ly α forest data and BAO, we found interesting constraints with the parameters restricted to the intervals $0.19 \leq \Omega_m \leq 0.23$ and $0.53 \leq h \leq 0.82$ within a 68.3% confidence level. All these results are in agreement with recent measurements reported in the literature (Table 1); however, due to the limited sample and our poor knowl-

edge of the IGM, they are weaker. We expect to obtain better and more reliable constraints on the Hubble constant in the near future by using a bigger sample and having a deeper understanding of the IGM (preferably with independent estimates for Γ and T_0). This will make our technique a true complementary of other estimates, with the advantage of being independent of the cosmic distance ladder.

The authors are grateful to J. V. Cunha and J. F. Jesus for their helpful discussions. VCB is supported by CNPq and RNG is supported by CAPES. JASL is partially supported by CNPq and FAPESP (No. 04/13668-0).

REFERENCES

- Ade P. A. R. et al., 2014, *A&A*, 571, A16
Aghaee, A., Petitjean, P., Srianand, R., Stalin, C. S., & Guimarães, R. 2010, *Journal of Astrophysics and Astronomy*, 31, 59
Aracil B., Petitjean P., Pichon C., Bergeron J., 2004, *A&A*, 419, 811
Amanullah R. et al., 2010, *ApJ*, 716, 712
Bergeron J. et al., 2004, *The Messenger*, 118, 40
Blake C. et al., 2012, *MNRAS*, 425, 405
Bolton J. S., Haehnelt M. G., Viel M., Springel V., 2005, *MNRAS*, 357, 1178

- Burles S., Nollett K. M., Turner M., 2001, *ApJ*, 552, L1
- Busca N. G. et al., 2013, *A&A*, 552, A96
- Busti V. C., Clarkson C., Seikel M., 2014, *MNRAS*, 441, L11
- Carswell R. F., Whelan J., Smith M., Bokserberg A., Tytler D., 1982, *MNRAS*, 198, 91
- Chand H., Srianand R., Petitjean P., Aracil B., 2004, *A&A*, 417, 853
- Chen G., Ratra B., 2011, *PASP*, 123, 1127
- Croft R. A. C., Weinberg D. H., Katz N., Hernquist L., 1998, *ApJ*, 495, 44
- Croft R. A. C., Weinberg D. H., Pettini M., Hernquist L., Katz N., 1999, *ApJ*, 520, 1
- Cunha J. V., Marassi L., Lima J. A. S., 2007, *MNRAS*, 379, L1
- Eisenstein D. J. et al., 2005, *ApJ*, 633, 560
- Faucher-Giguère C.-A., Lidz A., Hernquist L., Zaldarriaga M., 2008, *ApJ*, 682, L9
- Faucher-Giguère C.-A., Lidz A., Hernquist L., Zaldarriaga M., 2008, *ApJ*, 688, 85
- Freedman W. L. et al., 2001, *ApJ*, 553, 47
- Freedman W. L., Madore B. F., 2010, *ARA&A*, 48, 673
- Freedman W. L. et al., 2012, *ApJ*, 758, 24
- Guimarães R., Petitjean P., Rollinde E., de Carvalho R. R., Djorgovski S. G., Srianand R., Aghaee A., Castro S., 2007, *MNRAS*, 377, 657
- Hinshaw G. et al., 2013, *ApJS*, 208, 19
- Hui L., Gnedin N. Y., 1997, *MNRAS*, 292, 27
- Hui L., 1999, *ApJ*, 516, 519
- Jackson N., 2007, *Living Rev. Relativ.*, 10, 4
- Jimenez R., Verdi L., Treu T., Stern D., 2003, *ApJ*, 593, 622
- Holanda R. F. L., Cunha J. V., Marassi L., Lima J. A. S., 2012, *J. Cosmology Astropart. Phys.*, 1202, 035
- Kirkman D. et al., 2005, *MNRAS*, 360, 1373
- Komatsu E. et al., 2011, *ApJS*, 192, 18
- Lima J. A. S., Jesus J. F., Cunha J. V., 2009, *ApJ*, 690, L85
- Mandelbaum R., McDonald P., Seljak U., Cen R., 2005, *MNRAS*, 344, 776
- Marra V., Amendola L., Sawicky I., Valkenburg W., 2013, *Phys. Rev. Lett.*, 110, 241305
- McDonald P., Miralda-Escudé J., Rauch M., Sargent W. L. W., Barlow T. A., Cen R., Ostriker J. P., 2000, *ApJ*, 543, 1
- McDonald P., Miralda-Escudé J., 2001, *ApJ*, 549, L11
- McDonald P. et al., 2005, *ApJ*, 635, 761
- McQuinn M., Lidz A., Zaldarriaga M., Hernquist L., Hopkins P. F., Dutta S., Faucher-Giguère, C.-A., 2009, *ApJ*, 694, 842
- Meiksin A., White M., 2004, *MNRAS*, 350, 1107
- Miralda-Escudé J., Cen R., Ostriker J. P., Rauch M., 1996, *ApJ*, 471, 582
- Miralda-Escudé J., Haehnelt M., Rees M. J., 2000, *ApJ*, 530, 1
- Murdoch H. S., Hunstead R. W., Pettini M., Blades J. C., 1986, *ApJ*, 309, 19
- Nusser A., Haehnelt M., 1999, *MNRAS*, 303, 179
- Padmanabhan T., 2002, *Theoretical Astrophysics. Volume III, Galaxies and Cosmology*. Cambridge Univ. Press, Cambridge
- Paraficz D., Hjorth J., 2010, *ApJ*, 712, 1378
- Peebles P. J. E., 1993, *Principles of Physical Cosmology*. Princeton Univ. Press, Princeton
- Perlmutter S. et al., 1999, *ApJ*, 517, 565
- Pettini M., Zych B. J., Murphy M. T., Lewis A., Steidel C. C., 2008, *MNRAS*, 391, 1499
- Prochaska J. X., Gawiser E., Wolfe A. M., Castro S., Djorgovski S. G., 2003, *ApJ*, 595, 9
- Prochaska J. X., Castro S., Djorgovski S. G., 2003, *ApJS*, 148, 317
- Rauch M. et al., 1997, *ApJ*, 489, 7
- Riess A. G. et al., 1998, *AJ*, 116, 1009
- Riess A. G. et al., 2009, *ApJ*, 699, 539
- Riess A. G. et al., 2011, *ApJ*, 730, 119
- Rollinde E., Srianand R., Theuns T., Petitjean P., Chand H., 2005, *MNRAS*, 361, 1015
- Rollinde E., Theuns T., Schaye J., Pâris I., Petitjean P., 2013, *MNRAS*, 428, 540
- Sandage A., Tammann G. A., Saha A., Reindl B., Macchetto F. D., Panagia N., 2006, *ApJ*, 653, 843
- Sheinis A. I., Bolte M., Epps H. W., Kibric R. I., Miller J. S., Radovan M. V., Bigelow B. C., Sutin B. M., 2002, *PASP*, 114, 851
- Simha V., Steigman G., 2008, *J. Cosmology Astropart. Phys.*, 0806, 16
- Slosar A. et al., 2013, *J. Cosmology Astropart. Phys.*, 1304, 026
- Tammann G. A., Reindl B., 2013, *A&A*, 549, 136
- Tytler D. et al., 2004, *ApJ*, 617, 1
- Viel M., Bolton J.S., Haehnelt M.G., 2009, *MNRAS*, 399, L39
- Weinberg D. H., Davé R., Katz N., Kollmeier J.A., 2003, in Holt S. S., Reynolds C. S., eds, *AIP Conf. Proc. Vol. 666, The Emergence of Cosmic Structure: Thirteenth Astrophysics Conference*. Am. Inst. Phys., New York, p. 157
- Wyithe J. S. B., Bolton J. S., Haehnelt M. G., 2008, *MNRAS*, 383, 691
- Zaldarriaga M., Hui L., Tegmark M., 2001, *ApJ*, 557, 519

V. C. Busti: Astrophysics, Cosmology & Gravity Centre (ACGC), and Department of Mathematics & Applied Mathematics, University of Cape Town, Rondebosch 7701, Cape Town, South Africa.

R. N. Guimarães: CAPES Foundation, Ministry of Education of Brazil, Brasília, DF, Brazil (rguimara@on.br).

J. A. S. Lima: Instituto de Astronomia, Geofísica e Ciências Atmosféricas (IAG-USP), Universidade de São Paulo, 05508-900 São Paulo, SP, Brazil.