

WAVE PROPAGATION INTO THE GALACTIC ATMOSPHERE

M.A. Martos¹, D.P. Cox², and M. Walters²

We present hydrodynamical modelling of wave propagation into the Galactic atmosphere. Spectra consistent with observations are shown to be a natural consequence of the dynamical response of the thick Galactic disk of gas, cosmic rays and magnetic field to disturbances produced at the midplane of the Galaxy. The response to the spiral density wave is studied. Regardless of the source of the disturbance, an initial velocity gradient along the line of sight towards the Galactic halo has as a response the expansion of the disk followed by an eventual differential fall back of the structure, with the lower parts of the disk falling first and the outer parts later. The two regions are separated by a shock front that rapidly decelerates the infalling outer gas. The spectra so obtained reproduce naturally the main features of the kinematics observed above the Galactic plane, namely the two velocity components at roughly 0 and -50 km/s from the H I profile, and *HST* ultraviolet observations of the Galactic halo star HD 93521 presented by Spitzer & Fitzpatrick (1993).

¹Instituto de Astronomía, Universidad Nacional Autónoma de México

²University of Wisconsin-Madison, Dept. of Physics, USA

CONSTRAINTS ON THE ORIGIN OF ULTRA-HIGH ENERGY COSMIC RAYS

G.A. Medina Tanco¹, E.M. de Gouveia Dal Pino¹, and J.E. Horvath¹

We report the results of 3-D simulations of non-diffusive propagation of Ultra-High Energy Cosmic Rays (UHECR) ($E > 10^{20}$ eV) through the intergalactic and extended halo media. We quantify the expected angular and temporal correlations between the events and the sources for these distance scales. Our results indicate that the Yakuszk and Fly's Eye events probably lie outside the supergalactic plane. The so-proposed angular correlations with several extragalactic radio sources are found to be unlikely. The calculated time delays between UHE protons and gamma-ray counterparts do not match the claimed GRB-UHECR associations for either nearby extragalactic or extended halo distance scales. The present results impose rather severe constraints on the search of UHE acceleration sources.

¹Instituto Astronomômico e Geofísico, Universidade de São Paulo, Brazil

A SEARCH FOR SPECTROSCOPIC BINARIES IN THE MAGELLANIC CLOUDS

Virpi S. Niemela^{1,2,3}, Nidia I. Morrell^{1,4,5}
Guillermo L. Bosch^{1,5,6}, and Rodolfo H. Barbá^{1,4,5}

As part of an ongoing observing program devoted to the study of massive binary stars in the Magellanic Clouds, we have determined radial velocities for OB stars searching for variations due to binary motion. Based on spectroscopic observations obtained at CTIO, Chile, and CASLEO, Argentina, we present here the results of our study of the radial velocity behavior of 2 stars in the Small Magellanic Cloud (SMC), and 5 stars in the Large Magellanic Cloud (LMC).

In the SMC we performed a new determination of the radial velocity orbit of the WN3 + O binary AzV 332. We find that the absorption lines in this system arise from at least two different stars, thus indicating that AzV 332 is a multiple system.

In the SMC we also studied the B1Ia type star AzV 78, which turns out to have constant radial velocity within the observational errors. Our radial velocity values for this star are in good agreement with previously determined values.

We have found two new binaries among the stars studied in the LMC, namely Sk $-66^{\circ} 35$ (B1.5Ia) and Sk $-67^{\circ} 167$ (O4If). The former appears to be a long period single-lined binary with an elliptical orbit; while the latter shows double lines in its spectrum.

Marginal evidence of radial velocity variations was found from 8 spectra of Sk $-67^{\circ} 211$ (O3III).

No significant radial velocity variations were found in the available spectroscopic observations of Sk $-71^{\circ} 42$ (B2Ia) and Sk $-67^{\circ} 166$ (O4If).

¹Facultad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata, Argentina

²Member of Carrera del Investigador, CIC, Prov. de Buenos Aires, Argentina

³Visiting Astronomer CTIO, NOAO, operated by AURA for the NSF, USA

⁴Member of Carrera del Investigador, CONICET, Argentina

⁵Visiting Astronomer CASLEO, San Juan, Argentina

⁶Fellow of CONICET, Argentina