

INTERACTING ELLIPTICAL GALAXIES

Amelia Ramírez¹

Interacting galaxies show useful properties to constrain models of dark matter haloes. In general, these haloes seem to be as big as $200 \text{ h}^{-1} \text{ kpc}$ of radius in elliptical galaxies (Bahcall & Tremaine 1981; Charlton & Salpeter 1991; Zaritzky et al. 1993; Bahcall et al. 1995). But, we have a very poor information on the density profiles which determine the gravitational potential of those objects.

The dumbbell galaxies, two ellipticals of similar luminosities sharing a common luminous halo, are simpler than spiral pairs to study using N-body simulations. Since ellipticals have simpler morphological and internal dynamics the simulations produce less complicate distortions to be compared with observations. However, to perform these simulations it is required to know the initial condition of the encounter, the kind of distortion to be reproduced and take into account possible differences due the environment where the pair is located.

We present in this work results of a statistical, spectroscopic and photometric study of a sample of dumbbell galaxies in clusters (38% of the sample), groups (50%) and isolated (12%) (Ramírez et al. 1994). The dynamical analysis of the environment of one of these dumbbell pairs (NGC 4782/3) is in complete agreement with the existence of an extensive dark halo and the presence of a satellite system acting as test particles in the halo potential (Quintana et al. 1996). The analysis of the photometric data on a sub-sample shows pairs with distortions as offcentering isophotes and haloes with tails (18% of the sample) and bridges (56%), produced by tidal forces. In four pairs we found signs of a stellar population between the two components, possible due to new star formation region or exchange of stellar material (Ramírez & de Souza 1995).

- Bahcall, N.A., Lubin, L.M., & Dorman, V. 1995, ApJ 447, L81
 Bahcall, J.N., & Tremaine, S. 1981, ApJ 244, 805
 Charlton, J.C., & Salpeter, E.E. 1991, ApJ 375, 517
 Quintana, H., Ramírez, A., & Way, M.J. 1996, AJ 111, 603
 Ramírez, A., Hertling, G., & Quintana, H. 1994, Boll.Soc.Astr.Bras. 14, 82
 Ramírez, A., & de Souza, R. 1995, Boll.Soc. Astr.Bras. 15, 14
 Zaritzky, D., Smith, R., Frenk, C., & White, S.D.M. 1993, ApJ 405, 464

¹Depto. de Astronomia, IAG, Universidade de São-Paulo, Brazil

DISTRIBUTION OF INTERACTING GALAXIES IN A TEST SAMPLE OF CLUSTER OF GALAXIES

Carmen Gloria Ríos¹ and M. Faundez-Abans²

We present a study of the distribution of interacting galaxies in a test sample of clusters of galaxies. We propose a morphologic identification of the phenomena of interaction according to the visual appearance in the J plates of the ESO/SERC Southern Sky Survey. The test sample has 19 clusters of galaxies belonging to the South Hemisphere ($\delta < -16^\circ$), classified according to the Bautz-Morgan classification scheme. We studied the radial profile of the distribution of interacting galaxies to establish if the interaction has an intrinsic behavior related to the dynamic of the cluster more than a simple interaction between galaxies.

¹Departamento de Astronomía y Astrofísica, Pontificia Universidad Católica, Chile

²Laboratorio Nacional de Astrofísica, Brazil

NARROW-BAND IMAGERY OF NGC 4736

Irapuan Rodrigues¹, Horacio Dottori¹, Jordi Cepa², and José Vilchez²

In this work we study the galaxy NGC4736, using narrow band interference filters imaging centered at the emission lines $[\text{O II}]_{\lambda\lambda 3727+3729}$, $\text{H}\beta$, $[\text{O III}]_{\lambda 5007}$, $\text{H}\alpha$, $[\text{S II}]_{\lambda 6716+6730}$ and $[\text{S III}]_{\lambda 9070}$ and nearby continua at $\lambda 3812 \text{ \AA}$, $\lambda 4556 \text{ \AA}$, $\lambda 5276 \text{ \AA}$, $\lambda 6269 \text{ \AA}$, and $\lambda 9193 \text{ \AA}$. We used a CCD camera attached to the Cassegrain f/15 focus of the 1.0-m Jacobus Kapteyn Telescope, in the Roque de los Muchachos Observatory, La Palma, Canary Islands. The detector used was a GEC CCD 400×590 pixels (1 pixel = $22 \mu\text{m}$).

We have obtained sizes, positions, emission line absolute fluxes, and continua intensities for 90 H II regions, mainly distributed in a ring-like structure of 3.2 kpc in diameter. The $\text{H}\alpha$ luminosities are in the range $37.3 \leq \log L_{\text{H}\alpha} \leq 39.4 \text{ erg s}^{-1}$. H II region diameters were measured in $\text{H}\alpha$ image after a seeing deconvolution procedure. We obtained core and halo diameters (D_c and D_{100} , respectively) for each H II region. The H II regions size distribution presents a characteristic diameter $D_o = 115 \text{ pc}$ and follows

¹Instituto de Física, Universidade Federal do Rio Grande do Sul, Brazil

²Instituto Astrofísico de Canarias, Islas Canarias, Spain