

## NIR IMAGING SPECTROSCOPY OF IRAS F10214+4724: EVIDENCE FOR A STARBURST REGION AROUND AN AGN AT $z = 2.3$

H. Kroker<sup>1</sup>, R. Genzel<sup>1</sup>, A. Krabbe<sup>1</sup>, L.E. Tacconi-Garman<sup>1</sup>, M. Tecza<sup>1</sup>, and N. Thatte<sup>1</sup>

### RESUMEN

Presentamos espectroscopía en la banda  $K$ , con resolución de  $1''$ , de la galaxia IRAS F10214+4724, con  $z = 2.284$ . Encontramos que las líneas de emisión  $H\alpha$  y  $[N II]$  tienen diferentes distribuciones espaciales. Detectamos una componente ancha ( $\Delta v_{FWZP} \approx 3500 \text{ km s}^{-1}$  en  $H\alpha$ ). Concluimos que F10214+4724 es una galaxia muy luminosa, con un núcleo activo y un disco de formación estelar y además está amplificada por una lente gravitacional. Tanto el núcleo activo como el disco de formación estelar, contribuyen significativamente a la luminosidad total de  $\sim 10^{13} L_{\odot}$ .

### ABSTRACT

We report  $1''$   $K$ -band imaging spectroscopy of the  $z=2.284$  galaxy IRAS F10214+4724. We find that the rest-frame  $H\alpha$  and  $[N II]$  emission have different spatial extents. Furthermore, we detect broad ( $\Delta v_{FWZP} \approx 3500 \text{ km s}^{-1}$ )  $H\alpha$  emission. We conclude that F10214+4724 is a very luminous gravitationally lensed galaxy, which intrinsically contains both a type 1 AGN and a more extended star-forming disk. The AGN and circumnuclear star formation both contribute significantly to the total luminosity of  $\sim 10^{13} L_{\odot}$ .

*Key words:* GALAXIES: INDIVIDUAL: IRAS F10214+4724 — GALAXIES: STARBURSTS — INFRARED: GALAXIES

### 1. INTRODUCTION

The  $z=2.284$  IRAS source F10214+4724 has been interpreted to be the most luminous galaxy in the universe ( $\geq 10^{14} L_{\odot}$  for  $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$  and  $q_0 = 0.5$ ;  $1''0 = 5.3 \text{ kpc}$ ; e.g., Rowan-Robinson et al. 1991, 1993) being in an early superstarburst. The rest frame UV/visible emission lines resemble those of Seyfert II nuclei (e.g., Elston et al. 1994). However, recent imaging (Eisenhardt et al. 1995; Broadhurst & Lehár 1995; Graham & Liu 1995) strongly suggest magnifications between 5 and 100 due to a gravitational lens.

We observed F10214+4724 in the  $K$ -band with the MPE imaging spectrometer,  $3D$ , at the 3.5-m telescope on Calar Alto, Spain (see Weitzel et al. 1996). The total on-source integration time was 9000 s, the spectral resolution was  $R=500$  and the sampling  $0''.5$  pixel. The seeing during the observations was  $1''$  FWHM at  $2 \mu\text{m}$ .

### 2. RESULTS AND DISCUSSION

*Line and Continuum Maps: Differences in the Spatial Distribution.* Figure 1 shows line and continuum maps obtained with  $3D$  overlaid on the  $K$ -band image of Graham & Liu (1995; gray scale). Our key findings are the different spatial morphologies in all of these maps:  $H\alpha$  and  $[N II]$  line emission is restricted to the immediate neighborhood of the brightest source of the F10214+4724 complex (source 1 in Matthews et al. 1994). No significant line emission is found in sources 2, 3, or 4, anywhere in the  $K$ -band. The  $[N II]$  emission is fairly compact and marginally extended (east-west), as concluded by Matthews et al. (1994).  $H\alpha$  is also compact, but is significantly more extended east-west, following approximately the  $1''.5 - 2''$  arc structure of source 1 in the Keck  $K$ -band image. Line-free  $K$ -continuum is visible in source 1 as well, and has an extension north of it at the position of source 2 (Fig. 1, right panel). Source 2 is weaker than source 1 by a factor of 2. The integrated  $1.95 - 2.4 \mu\text{m}$  map (not shown) is in excellent agreement with the Keck  $K$ -band image. Thus, we conclude that a significant fraction of the east-west extent in the  $K$ -band maps is due to line emission.

<sup>1</sup>Max-Planck-Institut für extraterrestrische Physik (MPE), Postfach 1603, D-85740 Garching bei München, Germany.




Fig. 1. 3D-channel maps (contours; linear scale) of three spectral regions on top of the Keck  $K$ -band map of Graham & Liu (1995) (logarithmic grey scale). *left*:  $H\alpha$ . *middle*:  $[\text{N II}]\lambda 6583$ . Contours for both maps in  $2\sigma$  steps, starting with  $3\sigma$ . *right*: line-free  $K$ -continuum (98 continuum channels between  $2.02$  and  $2.32 \mu\text{m}$ ). Contours are in  $1\sigma$  steps, starting with  $1\sigma$ .

**Broad  $H\alpha$  Emission.** We have fitted multiple Gaussians to the  $H\alpha/[\text{N II}]\lambda\lambda 6548, 6583$  spectra for different spatial regions: aperture sizes of  $0''.75, 1''.25, 1''.75$  and  $2''.25$ , centered on the  $H\alpha$  peak; two ringlike regions containing the flux from the  $2''.25$  aperture, but not the central  $0''.75$  and  $1''.25$  aperture, respectively; and, finally, a region covering just the wings (more than  $0''.7$  east-west distance from the center) of the  $2''$  arc (source 1). Fixing the  $[\text{N II}]\lambda 6583/[\text{N II}]\lambda 6548$  ratio to its theoretical value, results in a fit with unacceptably large residuals (Fig. 2a) for a fit with three Gaussians. It is necessary to add emission contributing blueward of the  $H\alpha$  peak to account for the line wing to the blue of  $[\text{N II}]\lambda 6548$ . The most plausible such component is a broad  $H\alpha$  emission line. As demonstrated in Figure 2 b-d, the fits using a fourth, broad Gaussian centered on the narrow  $H\alpha$  line are very satisfactory. The contribution of broad  $H\alpha$  to the total line flux of the  $H\alpha+[\text{N II}]$  line system is 50% for the  $2''.25$  aperture, the line width is  $2400 \text{ km s}^{-1}$  FWHM (FWZP= $3500 \text{ km s}^{-1}$ ) for all fitted regions, while the three narrow lines are unresolved (FWHM  $600 \pm 100 \text{ km s}^{-1}$ ). Thus, we conclude that the nucleus of F10214+4724 contains a buried BLR and, therefore, is likely to have a type 1 (Seyfert I or QSO) nucleus.

**Extended  $H\alpha$  Emission: Circumnuclear Star Formation.** The  $[\text{N II}]\lambda 6583/H\alpha_{\text{broad}}$  line ratio does not vary with aperture size, which is consistent with a dominating compact AGN origin of both lines. Therefore, we finally constrained the fit by fixing the  $[\text{N II}]\lambda 6583/H\alpha_{\text{broad}}$  line ratio and fitted six parameters to our data set: a narrow and a broad linewidth, two amplitudes, a wavelength position and a continuum level. In contrast, the  $[\text{N II}]\lambda 6583/H\alpha_{\text{narrow}}$  line ratio shows spatial variation in our data; it drops from  $1.66 \pm 0.06(1\sigma)$  in the peak-centered  $0''.75$  aperture to  $1.2 \pm 0.16$  within the  $2''.25$  aperture. In the extreme wings of the  $2''$  arc the ratio is  $0.6 \pm 0.2$ . Thus, we find that 30% of the total narrow  $H\alpha$  flux is in an east-west extended component, corresponding to the  $2''$  arc in the Keck  $K$ -band image. For small apertures, the  $[\text{N II}]\lambda 6583/H\alpha_{\text{narrow}}$  line ratio is consistent with a classical photoionized NLR. In the east-west wings that undiluted ratio is  $\leq 0.5$  and resembles that of an HII region galaxy ( $[\text{N II}]\lambda 6583/H\alpha=0.2-0.5$ , Osterbrock 1989, p. 346). We propose that this extended component is a starforming disk and may be associated with the zone of FIR continuum and submm CO emission. Although the S/N substantially decreases in the outer region, the main signature of the changing physical conditions, which is the change in  $[\text{N II}]/H\alpha$  ratio, is a robust result. Even the fourth component ( $H\alpha_{\text{broad}}$ ) is not critical at all for this conclusion; in fact just fitting three components without fixing the  $[\text{N II}]\lambda 6583/[\text{N II}]\lambda 6548$  flux ratio would make the result even clearer because the  $[\text{N II}]\lambda 6583/H\alpha_{\text{narrow}}$  line ratio would drop even more dramatically in the outer regions.

**Energetics of F10214+4724: Contributions of AGN and Star Formation.** From the observed broad- and narrow-line  $H\alpha$  fluxes, estimates of their extinctions, and model gravitational lens magnifications, it is now possible to derive a first rough estimate of the luminosities of these two components. Lens models (Broadhurst & Lehár 1995; Eisenhardt et al. 1995) predict a magnification of 50–100 for the central ( $0''.7$ , intrinsic size




Fig. 2. Line fits to the  $H\alpha$ + $[N\ II]\lambda\lambda$  6548, 6583 profile for different apertures: (a) :  $0.''75$ , three Gaussians; (b-d) :  $0.''75$ ,  $1.''75$ , and Annulus ( $2.''25 - 0.''75$ ), fitted by four Gaussians. Histogram represents the data, short-dashed lines the single Gaussian fits, and long-dashed line the sum. Lower plots are the residuals of the fits.

TABLE 1  
DERIVED LUMINOSITIES OF VARIOUS COMPONENTS

Component	Observed Luminosity <sup>a</sup>	Extinction Correction	Magnification Factor	$L_{bol}/L_{H\alpha}$	$L_{bol}$
$H\alpha_{broad}$	$16 \pm 4$	8	50-100	400-600 <sup>b</sup>	500-1500
$H\alpha_{narrow}$	$5 \pm 1$	8	50-100	180 <sup>b</sup>	70-140
$H\alpha_{starformation}$	$1.5 \pm 0.5$	10-100	5-10	200 <sup>c</sup>	300-6000

<sup>a</sup>All luminosities are in  $10^9 L_{\odot}$ .

<sup>b</sup>Netzer 1990.

<sup>c</sup>case B recombination (e.g., Osterbrock 1989) and  $L_{bol}/L_{Lyc} = 8$ .

$\sim 0.01''$  or 50 pc), bright component. For the more extended ( $2''$ ,  $0.4''$  or 2 kpc intrinsic), fainter arc visible on the Keck-image, the magnification is 5-10. This is probably also the size scale of the FIR continuum and CO emission region (Downes et al. 1995). In Table 1 we give an estimate of the BLR, NLR and starburst contributions to the  $H\alpha$  flux. The visual extinction can be derived from the observed  $H\alpha/H\beta$  ratio of 8.6 (Iwamuro et al. 1995). For an intrinsic ratio of 3.1 this results in  $A_V = 2.9$  and an extinction correction of a factor of 8. For the circumnuclear region a mixed dust-gas model is more likely, leading to much larger values for  $A_V$ . The CO fluxes of Downes et al. (1995) convert to  $A_V \geq 10^2$  in the circumnuclear star forming region and an extinction correction of at least 10 or even  $10^2$ . Finally, we estimated the  $L_{H\alpha}$  to  $L_{bol}$  (case B recombination) for the star forming region, and for typical AGN NLR/BLR models (Netzer 1990) (see Table 1). While the details are still quite uncertain, the basic result is clear. Both a powerful AGN and a very luminous circumnuclear star formation region contribute to the total luminosity of F10214+4724. Thus, F10214+4724 has intrinsically similar characteristics as other luminous IRAS galaxies, although it is still one of the most luminous galaxies.

#### REFERENCES

- Broadhurst, T., & Lehar, J. 1995, ApJ, 450, L41  
 Downes, D., Solomon, P. M., & Radford, S. J. E. 1995, ApJ, 453, L65  
 Elston, R., et al. 1994, AJ, 107, 910  
 Eisenhardt, P. R., Armus, L., Hogg, D. W., Soifer, B. T., Neugebauer, G., & Werner, M. W. 1996, ApJ, 461, 72  
 Graham, J. R., & Liu, M. C. 1995, ApJ, 449, L29  
 Iwamuro, F., Maihara, T., Tsukamoto, H., Oya, S., Hall, D. N. B., & Cowie, L. L. 1995, PASJ, 47, 265

- 1997RMxAC...6...48K
- Kroker, H., Genzel, R., Krabbe, A., Tacconi-Garman, L. E., Tecza, M., & Thatte, N. 1996, ApJ, 463, L55  
Matthews, K., et al. 1994, ApJ, 420, L13  
Netzer, H. 1990, in Active Galactic Nuclei, Saas-Fee Advanced Course 20 Lecture Notes, ed. T. J.-L. Courvoisier & M. Mayor (Berlin: Springer), 93  
Osterbrock, D. E. 1989, Astrophysics of Gaseous Nebulae and Active Galactic Nuclei, University Science Books  
Rowan-Robinson, M. et al. 1991, Nature, 351, 719  
Rowan-Robinson, M. et al. 1993, MNRAS, 261, 513  
Weitzel, L., Krabbe, A., Kroker, H., Thatte, N., Tacconi-Garman, L. E., Cameron, M., & Genzel, R. 1996, A&A, in press