

ABSTRACTS OF POSTERS

CURRENTS, FIELDS, AND POTENTIALS
IN THE IONOSPHERE OF VENUSD. M. Hurley¹ and P. A. Cloutier¹

Law and Cloutier have developed a complete picture of the magnetic field in the dayside ionosphere of Venus. Their investigation is based upon observations from the Pioneer Venus Orbiter (PVO) magnetometer instrument. We embed the model magnetic field to a model ionosphere and perform several analyses. First of all, we determine the electric currents associated with the magnetic field configuration. We use the model atmosphere to transform the current into an electric field map of the region. Then, we examine the field aligned current and cross field current to investigate the potential drops along field lines. In doing so, we derive the map of the ionospheric electric potential for the dayside of Venus.

¹ Dept. of Space Physics and Astronomy, Rice University, Houston, TX, USA; dana@spacsun.rice.edu.

SPECTRA AND NEURAL NETWORKS FOR
THE SPECIFICATION OF STORM-TIME
ENERGETIC ELECTRONS AT
GEOSTATIONARY ORBITJ. W. Freeman¹ and T. P. O'Brien¹

Electrons whose energy is of the order MeV have been implicated in the failure and malfunction of Geostationary spacecraft. It is, therefore, important to be able to specify and even forecast the flux of these particles during and following geomagnetic storms. A first step is the understanding of their spectral properties and their relationship to lower energy electrons that can already be modeled.

At Geosynchronous orbit, the fluxes of electrons whose energies range from 100 keV to 2 MeV fall off with increasing energy according to a power-law. On a log-log plot, this spectrum appears as a line, with

slope and offset parameters. We note that the spectrum parameters vary in time, but that the overall power-law persists. We present our preliminary efforts modeling these parameters with an Artificial Neural Network. We have begun our investigation with the geomagnetic storm in early November, 1993.

We report initial success using simultaneous input values from Dst, satellite local time, and satellite-measured flux in the 30–45 keV channel. The ANN performs reasonably well throughout most of the storm and during the recovery phase. However, we believe the simultaneous input to be insufficient for modeling the full behavior of the spectrum. We have not yet implemented any historical information as inputs to the ANN. With such additional inputs, we expect the accuracy to improve.

¹ Dept. of Space Physics and Astronomy, Rice University, Houston, TX, USA; freeman@alfven.rice.edu.

SPECTROPHOTOMETRIC DATA OF THE
CENTRAL STAR OF THE PLANETARY
NEBULA LMC N66. QUANTITATIVE ANALYSIS
OF ITS WN TYPE SPECTRUM¹M. Peña², W.-R. Hamann³, L. Koesterke³,
J. Maza⁴, R. H. Méndez⁵, M. Peimbert²,
M. T. Ruiz⁴, and S. Torres-Peimbert²

HST, *IUE* and ground-based observations of the central star of the LMC planetary nebula N66 (CS N66), obtained in different epochs, are presented. Since 1990 CS N66 has shown remarkable short- and long-term spectroscopic and photometric changes amounting to more than 3 magnitudes in the optical. Expanding model atmospheres have been constructed to fit observations from different epochs. Fits provide the chemical composition, the fundamental stellar parameters, L_* , T_* , R_* , the mass-loss rate and the wind velocity. From our best models we found that CS N66 is a very luminous He star ($X/Y \leq 0.1$), with a small amount of N, under-

going a violent and unstable mass loss event. The photospheric chemical abundances correspond to the equilibrium CNO nuclear burning values, while the nebula has a normal chemical composition. Models fitting data from different epochs show that the fundamental stellar parameters remain constant with time, with values $\log L_*/L_\odot = 4.53 \pm 0.10$, $T_* = 93\,300$ K, and $R_* = 0.71 R_\odot$. The short- and long-term stellar variations are produced by large changes in the mass-loss rate, which varies by large factors, from $\dot{M} \leq 8 \times 10^{-7} M_\odot \text{ yr}^{-1}$ in 1983 (pre-outburst epoch) to $\dot{M} = 2.5 \times 10^{-5} M_\odot \text{ yr}^{-1}$ in early 1995 (maximum stellar brightness). No evidence to support the suggestion that the outburst was due to a late thermal pulse was found. We propose that the event taking place in CS N66 was produced by an atmospheric instability similar to those triggering the giant eruptions of Population I LBV stars. We briefly discuss an approach to the Eddington-limit due to changes in the opacity and a non-radial pulsational instability as two possible mechanism which could have caused the atmospheric outburst.

¹ Based on observations made with the NASA/ESA *Hubble Space Telescope*, the *IUE* satellite, and at Cerro Tololo InterAmerican Observatory.

² Instituto de Astronomía, UNAM, México, D.F., México; miriam@astroscu.unam.mx.

³ Lehrstuhl Astrophysik der Universität Potsdam, Am Neuen Palais 10, D-14469 Potsdam, Germany.

⁴ Depto. de Astronomía, Universidad de Chile, Santiago, Chile.

⁵ Institut für Astronomie und Astrophysik, Universitäts-Sternwarte München, D-81679, München, Germany.

SHAPING BIPOLAR AND ELLIPTICAL PLANETARY NEBULAE: EFFECTS OF STELLAR ROTATION AND MAGNETIC FIELDS

G. García-Segura¹, N. Langer²,
M. Różyczka³, and J. Franco¹

We present hydrodynamical and magneto-hydrodynamical simulations for the evolution of planetary nebulae formed through the interaction of two succeeding constant stellar winds. The latitudinal dependence of the wind parameters are described by a single wind function in our calculations. We explore various relevant parameters, including the effects of stellar rotation, ionizing radiation field and stellar magnetic field, and a catalogue of resulting nebula shapes is generated.

We find that strongly bipolar nebulae can be obtained when the stellar rotational velocity, v_{rot} , approaches the critical rotational velocity, v_{crit} , at the asymptotic giant branch (AGB) phase. As an exam-

ple, the shape of the Hourglass Nebula (MyCn18) is obtained for the value of the ratio $\Omega = v_{\text{rot}}/v_{\text{crit}} \approx 0.98$. We estimate the evolution of the rotation rate and of the critical rotation rate in thermally pulsing AGB stars and find a plausible scenario which predicts critical rotation at the tip of the ABG for single stars with an initial mass of more than $\sim 1.3 M_\odot$.

When a stellar magnetic field is combined with the effect of the rotation at the AGB phase, highly collimated bipolar nebulae can be obtained such as M2-9 or He 2-437. Provided that the field is sufficiently strong, the formation of ansae and jets in the polar regions of the nebula have also been found, such as IC 4593. On the other hand, weaker fields can account for classical elliptical nebulae such as NGC 6905 even in nonrotating (i.e., spherically symmetric) AGB winds.

Photoionization was found to be important in producing dynamical effects. It generates irregularities in the shape of the simulated nebulae, and may be responsible for the formation of irregularly shaped planetary nebulae, such as Sh 2-71 or WeSb 4. It also leads to the formation of cometary knots, preferentially at the equatorial region, similar to those seen in the Helix nebula (NGC 7293).

¹ Instituto de Astronomía, UNAM, México D. F., México; ggs@astroscu.unam.mx.

² Institut für Theoretische Physik und Astrophysik, Universität Potsdam, D-14415 Postdam, Germany.

³ Warsaw University Observatory, Al. Ujazdowskie 4, 00-478 Warszawa, Poland.

CHEMODYNAMICAL MODEL OF THE GALAXY: ABUNDANCE GRADIENTS PREDICTED FOR H II REGIONS AND PLANETARY NEBULAE

C. Allen¹, L. Carigi¹, and M. Peimbert¹

We present a chemodynamical evolution model of the Galaxy to determine chemical abundance gradients of different stellar populations. From this model we have determined the abundance gradients expected for H II regions as well as for planetary nebulae of different ages and different kinematical properties. We have compared the model predicted gradients with those derived from PNe of Types I, II and III. From this comparison we conclude that only about half of the stars evolving toward the white dwarf stage produce PNe and that the less massive stars are less likely to produce PNe. Other arguments supporting the previous conclusions are presented.

¹ Instituto de Astronomía, UNAM, México, D.F., México; chris@astroscu.unam.mx.