

endpoints to star formation ridges and dust lanes in two-armed spirals. In a previous paper (Puerari & Dottori 1992), we proposed a method to determine the leading or trailing character of the density wave perturbation in spiral galaxies by analyzing the distribution of H II regions. We idealize a new method based on the Fourier analysis of azimuthal profiles, to locate the CR and determine the arm character (trailing or leading) in spiral galaxies. Basically, we compare the behavior of the phase angle of the two-armed spiral in blue and infrared colors that pick out, respectively, young and older disk stellar population. The full paper of this poster contribution was published elsewhere (Puerari & Dottori 1997) and the reader can get more information on that paper. With our new method we confirm for NGC 7479 the existence of the leading pattern with CR at the extreme of the bar. We found in this galaxy the existence of an internal CR, indicating a trailing pattern. NGC 1832 presents three CRs, the inner and the outer ones indicate leading pattern and the intermediate CR, a trailing one. The most plausible physical interpretation for this situation is the existence of two pattern speeds.

- Beckman, J. E., & Cepa, J. 1990, *A&A*, 229, 37  
 Elmegreen, B. G., Elmegreen, D. M., & Montenegro, L. 1992, *ApJS*, 79, 37  
 Puerari, I., & Dottori, H. 1992, *A&AS*, 93, 469  
 \_\_\_\_\_, 1997, *ApJ*, 476, L73  
 Schweizer, F. 1976, *ApJS*, 31, 313

<sup>1</sup> Instituto Nacional de Astrofísica, Óptica y Electrónica, Puebla, Pue., México; puerari@inaoep.mx.

<sup>2</sup> Departamento de Astronomia, Instituto de Física, UFRGS, Porto Alegre, Brazil; dottori@if.ufrgs.br.

of Hg at visible wavelengths. Corrector distortion has limited resolution to 0.95 arcsec FWHM. Atmospheric seeing at the telescope's location (National Solar Observatory, Sunspot, NM) yields a nominal resolution of 1.3 arcsec (") FWHM. After a corrector fix in late 1997, we expect to obtain 0.5" FWHM images with good seeing.

The LMT was constructed with the purpose of characterizing the orbital space debris environment for NASA and conducting very wide/deep field astronomical surveys. Project costs were 1.2 M\$ or 5% of the estimated cost for a conventional (glass/pointable) 3.0-m telescope. Since March 1996, we have obtained 100+ hrs of orbital debris observations with various image intensifiers. The present debris detection size is < 4 cm at 1000 km assuming a 0.1 albedo spherical reflector. For the astronomical survey, we are using a Ford 2K thick CCD with 15  $\mu$ m pixels yielding a 0.6"  $\text{pix}^{-1}$  plate scale. The CCD is used in Time Delay Integration (TDI) mode wherein we drift scan at the sidereal rate to yield a 97.0 sec exposure time over a 20 arcminute FOV. To date, we have obtained (BVRI) and narrowband (400–1000 nm; 10–30 BW) photometry over 40+ sq. degrees of sky at high galactic latitude centered on +33° Dec. The accumulating data set (200+GB) is being used primarily to generate a wedge diagram extending to a redshift of 0.5 over 10 hrs of Right Ascension and containing approximately 1.5 million galaxies/QSOs; the largest survey of its kind by any telescope to date.

<sup>1</sup> Project Scientist, NASA Orbital Debris Observatory.

<sup>2</sup> Dept. of Space Physics and Astronomy, Rice University, Houston, TX, USA; mulrooney@sn3.jsc.nasa.gov.

## NASA 3.0-M LIQUID MIRROR TELESCOPE

M. K. Mulrooney<sup>1,2</sup>

We have constructed a 3.0-meter aperture zenith-staring telescope of an unusual nature. This Liquid Mirror Telescope (LMT) has as its primary mirror a spinning container of elemental mercury (Hg). We rely on the well-known principle that the equilibrium surface configuration of a rotating fluid is a paraboloid (if the axis of rotation is parallel to a uniform gravitational field). In our configuration, the primary container supports a 1.8 mm film of Hg and rotates on an air bearing spindle at 10 rpm, yielding a focal length of 4.5 m ( $f/1.5$ ). A Wynne-type corrector removes aberrations and produces a 6 cm diameter, 0.64 deg focal plane at prime focus. Optical throughput is 65% allowing for the 78% reflectivity

## EVIDENCE FOR SATURATED INVERSE COMPTON SCATTERING IN GAMMA-RAY BURSTS

A. Crider<sup>1</sup>, E. Liang<sup>1</sup>, and I. Smith<sup>1</sup>

Early studies of gamma-ray bursts found that the low-energy (30–100 keV) asymptotic spectral slope was typically 0 to  $-1.5$  (Band et al. 1993). However, we examined the time-resolved low-energy GRB spectra measured with the Burst and Transient Source Experiment (BATSE) and found that the asymptotic power slope is often positive (as high as  $1.6 \pm 0.3$ ) near the beginning of a burst and becomes negative as the burst progresses (Crider et al. 1997). These findings rule out many proposed emission mechanisms but can naturally be explained with saturated Comptonization as an emission mechanism in GRBs. By running Monte Carlo simulations, we

find that a simple decrease in the Thompson depth is all that is required to explain the observed change in the low-energy slope (Liang et al. 1997). Further evidence may be provided by recently discovered counterparts to GRB970228, observed with the Italian X-ray satellite *BeppoSAX* (Costa et al. 1997) and *Hubble Space Telescope* (Sahu et al. 1997). These counterparts fade at a rate that is consistent with saturated Comptonization. If this is indeed the emission mechanism, these observations place limits on the rate at which the spectral break energy decreases.

Band, D. L. et al. 1993, ApJ, 413, 281  
 Crider, A., et al. 1997, ApJ, 479, L39  
 Liang, E., et al. 1997, ApJ, 479, L35  
 Costa, E. et al. 1997, IAU Circ. No. 6576  
 Sahu, K. C. et al. 1997, Nature, in press

<sup>1</sup> Dept. of Space Physics and Astronomy, Rice University, Houston, TX, USA; acrider@spacsun.rice.edu.

#### IS THE EGRET SOURCE 2EG2020+4026 A GEMINGA-LIKE PULSAR?

K. T. S. Brazier<sup>1</sup>, G. Kanbach<sup>2</sup>, A. Carramiñana<sup>3</sup>,  
 J. Guichard<sup>3</sup>, and O. Reimer<sup>2</sup>

The error box of the EGRET source 2EG2020+4026, inside the  $\gamma$ -Cygni supernova remnant, has been found to contain a point-like X-ray source, denoted

as RX J2020.2+4026. Through optical observations, we consider unlikely the association of the X-ray source with the only coincidental star, a  $m_v \approx 15$  star of type K0V with no signs of coronal activity. The most direct interpretation of the data available is that 2EG2020+4026 is a young pulsar: the  $\gamma$ -ray emission shows no time variability and the spectrum is similar to that of Geminga; there is an X-ray point source inside the error box and SNR, and the  $F_\gamma/F_X$  is higher than in all objects apart from pulsars; and the lack of a credible optical counterpart is also consistent with a  $\gamma$ -ray emitting neutron star.

The proof that 2EG2020+4026 is a pulsar would be the detection of pulsations. We searched for these in a sample of EGRET data corresponding to GCRO observation 203 containing 1129 photons events with  $E > 200$  MeV. The search comprise most of the range of timing parameters of known pulsars,  $\nu \lesssim 30$  Hz and  $\dot{\nu} \lesssim 2.2 \times 10^{-11} \text{ Hz}^{-2}$ , and no coherent pulsations of sufficient statistical significance were found after  $3.78 \times 10^9$  tries. Its sensitivity was hampered by the high background in the source region and it does not constrain the pulsed fraction of the  $\gamma$ -ray flux. A thorough search requires higher S/N X-ray or  $\gamma$ -ray data.

Brazier K. T. S. et al. 1996, MNRAS, 281, 1033

<sup>1</sup> Dept. of Physics, University of Durham, Durham DH1 3LE England; Karen.Brazier@durham.ac.uk.

<sup>2</sup> Max-Planck Institut für Extraterrestrische Physik, Germany.

<sup>3</sup> Astrofísica, INAOE, Tonantzintla, Pue., México.

#### LIST OF ABSTRACTS OF POSTERS

CURRENTS, FIELDS, AND POTENTIALS IN  
 THE IONOSPHERE OF VENUS *D. M. Hurley*  
 & *P. A. Cloutier* 215

SPECTRA AND NEURAL NETWORKS FOR  
 THE SPECIFICATION OF STORM-TIME ENER-  
 GETIC ELECTRONS AT GEOSTATIONARY OR-  
 BIT *J. W. Freeman* & *T. P. O'Brien* 215

SPECTROPHOTOMETRIC DATA OF THE CEN-  
 TRAL STAR OF THE PLANETARY NEBULA  
 LMC N66. QUANTITATIVE ANALYSIS OF ITS  
 WN TYPE SPECTRUM *M. Peña,*  
*W.-R. Hamann,* *L. Koesterke,* *J. Maza,*  
*R. H. Méndez,* *M. Peimbert,* *M. T. Ruiz,* &  
*S. Torres-Peimbert* 215