

STUDY OF THE SYMBIOTIC STAR AR PAVONIS

C. Quiroga¹, E. Brandi², O. Ferrer¹, L. García, R. Barbá¹

Facultad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata, Argentina

and

F. Scaltriti

Observatorio Astronómico de Torino

RESUMEN

Se presenta un estudio espectroscópico y polarimétrico de la binaria eclipsante AR Pavonis. Se analizaron algunos espectros reunidos entre los años 1990 y 1998, y datos polarimétricos correspondientes al intervalo 1995 - 1998. A partir de las variaciones espectroscópicas observadas a lo largo del período orbital se estudió la naturaleza del sistema. Estudios polarimétricos indican que AR Pav presenta polarización intrínseca.

ABSTRACT

We present a study of the optical spectra and linear polarization of the eclipsing binary AR Pavonis. The optical spectra were obtained between 1990 and 1998, and polarimetric data between 1995 and 1998. From spectroscopic variations along the orbital motion we analyzed the nature of the system. Polarimetric studies indicate that AR Pav has intrinsic polarization.

Key Words: **POLARIZATION — SYMBIOTIC STARS**

1. INTRODUCTION.

The symbiotic stars are interacting binary systems composed of a cool giant star and a hot component surrounded by an ionized nebula. AR Pav is an eclipsing symbiotic binary with an orbital period of 604.5 days (Bruch et al. 1994). Thackeray & Hutchings (1974) proposed a model consisting of a red giant filling its Roche lobe losing mass to an evolved hot component, for this star. Kenyon & Webbink (1984) replaced the white dwarf with an accretion disk around a main sequence star.

2. OBSERVATIONS.

Spectroscopic observations were performed with the 2.15 m telescope of CASLEO, San Juan, Argentina. Intermediate resolution spectra were taken with a Boller & Chivens Cassegrain spectrograph, using a Z-Machine detector and a Thomson CCD of 384×576 pixels. High resolution spectra were obtained using a REOSC spectrograph and a Tek CCD of 1024×1024 pixels. Polarimetric observations were obtained with the photopolarimeter of Torino attached to the 2.15 m telescope of CASLEO. This instrument allows simultaneous measurements of polarization in five colour bands UBVRI.

3. SPECTROSCOPIC FEATURES.

The spectrum of AR Pav shows emission lines of H I, He I, He II $\lambda 4686$, N III, C III, Fe II, Ti II, [O III], [Fe II], [O I], and [O II] and molecular bands of TiO. The emission lines of H β , H γ , and He II $\lambda 4686$ have central

¹Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina.

²Comisión de Investigaciones Científicas de la Provincia de Buenos Aires (CIC), Argentina.




Fig. 1. 1a (*Left*): Radial velocities of H β and He II λ 4686. Filled circles correspond to radial velocities of all profile, open triangles, circles and cross correspond to velocities of red peak, central absorption and blue peak, respectively. 1b (*Right*): Percentage of polarization and position angle as a function of orbital phase for the filters UBVI.

absorption cores, also present in some emission lines of He I and [O III] λ 5007. The absence of emission in [Ne V], [Fe VI] and [Fe VII] lines, and of two broad Raman emissions $\lambda\lambda$ 6825, 7082, indicates a low ionization temperature for AR Pav.

Figure 1a shows the behavior of radial velocities of H β and He II λ 4686 emission lines with the orbital phase (ephemeris Bruch et al. 1994). Both radial velocities follow the orbital motion of the hot component. The same behavior is observed for the central absorption of H β , contrary to the observations of Thackeray & Hutchings (1974), who assume that the central absorption is originated in an expanding shell surrounding the whole system.

4. POLARIMETRY

Figure 1b shows the percentage of polarization and the position angle as a function of orbital phase for each filter UBVI. The percentage of polarization varies strongly at filters U, B, and V, reaching the maximum at phase 0.54 and 0.93. At these phases the percentage of polarization suggests Mie scattering (λ^{-1} dependence), which is produced by large particles. Large rotations of position angle are observed at phase 0.21 and 0.70 for filters B, V, R, and I.

5. CONCLUSIONS

The emission lines of H I and He II λ 4686 in AR Pav follow the orbital motion of the hot component. The central absorption present in these emission lines are most probably formed in a disk surrounding the hot component. The behavior of polarization indicates intrinsic polarization of the system, which is related with the orbital phase.

REFERENCES

- Bruch, A., Niehues, M., Jones, A. 1994, A&A, 287, 829
 Kenyon, S., & Webbink, R. 1984, ApJ, 279, 252
 Thackeray, A., & Hutchings, J. 1974, MNRAS, 167, 319



Focal Point 2b:

Compact Stars and Supernovae Progenitors



Dany Page and Andreas Reisenegger discussing about compact stars.