

## SEARCHING FOR MOVING GROUPS WITH *HIPPARCOS*

Luis Aguilar

Obs. Astronómico Nacional e Instituto de Astronomía, UNAM, Ensenada B. C., México

and

R. Hoogerwerf

Sterrewacht, Leiden, the Netherlands

### RESUMEN

Se presenta una búsqueda de grupos en movimiento en la vecindad solar. Se utiliza la base de datos de *HIPPARCOS* y un nuevo método de búsqueda llamado el *Método del Espaghetto*. Este método está optimizado para utilizar toda y únicamente la información astrométrica dada por *HIPPARCOS*: posición en el cielo, paralaje y movimiento propio. Hemos determinado la efectividad del método aplicándolo a realizaciones de Monte Carlo de grupos en movimiento artificiales. El método ha sido utilizado para encontrar grupos en movimiento en la vecindad solar

### ABSTRACT

We present a search for moving groups in the solar neighborhood. The search is done using the *HIPPARCOS* database and a new search method: *the Spaghetti Method*. This method is optimized to search using all, and only, the astrometric information provided by *HIPPARCOS*: position in the sky, parallax and proper motion. We have gauged the effectiveness of the method applying it to several Monte Carlo experiments with synthetic moving groups. This method has been used extensively to search for moving groups in the solar neighborhood.

*Key Words:* **ASTROMETRY: MOVING GROUPS**

The Hipparcos astrometric mission has provided us with a database of slightly more than  $10^5$  bright stars in the solar neighborhood, for which positions have been determined with a median uncertainty of 0.77 mas, parallaxes with a median error of 0.97 mas, and proper motions with a median precision of 0.88 mas/yr (ESA 1997). An important aspect of this database is that the astrometry is absolute all across the sky, thus opening up the possibility of searching for stars that share kinematics, regardless of their spread in the sky.

We have devised a maximum information method to search for moving groups in the *HIPPARCOS* database (Hoogerwerf & Aguilar 1999). The astrometry from *HIPPARCOS* restricts the tip of the velocity vector of a star to a line in velocity space parallel to the line of sight and offset from the origin by the implied tangential velocity vector. Any 3-dimensional velocity vector whose tip lies on this line will project on the plane of the sky to the measured tangential velocity vector. Errors in the measurement of the astrometric quantities will thicken this line into a cylinder of elliptical cross section (spaghetti).

Stars that belong to the same moving group, will share, within some intrinsic group velocity dispersion, the same 3-dimensional velocity vector and thus their respective spaghetti will intersect in velocity space. Since the probability of a chance intersection among random cylinders in 3-dimensional space quickly shrinks as the number of cylinders increases, any prominent intersection will be a candidate moving group. A series of random realizations of the null hypothesis of no moving groups is used to assess the statistical significance of the suspect intersections (Fig. 1).

The method has been compared to the traditional convergent point method (de Bruijne 1999), and with the results obtained by Perryman et al. (1998) (P98) for the Hyades. It has also been used extensively by de Zeeuw et al. (1999) in their extensive search for OB associations in the solar neighborhood. Finally, Aguilar




Fig. 1. The spaghetti Method: The first step is the selection of a volume in configuration space where a subsample is chosen according to user defined selection criteria. The spaghetti for each star is then computed and a Spaghetti Density Function ( $SDF$ ) is defined as the sum of the probability densities contributed by each spaghetti. Peaks in the  $SDF$  above a given threshold value  $SDF_0$  are then identified as suspect moving groups. Monte Carlo experiments with synthetic surveys with no moving groups are then used to assess the statistical significance of each suspect peak. Finally individual membership probabilities are computed.

& Hoogerwerf (1999) have used it to set upper limits on the velocity and number of members of putative halo moving groups passing within the solar neighborhood. In the case of the Hyades, the 3 components of the group velocity coincide within 0.3 km/s with those measured by P98, who had access to radial velocity information. Our method identifies 168 stars of which only 6 are not identified as Hyades members by P98.

L. A. thanks Sterrewacht Leiden for its hospitality. This research has been funded by NWO, the Netherlands Organization for Scientific Research, and in México by UNAM through DGAPA grant N114496 and by CONACyT through grant 27561-E. We also thank IAUNAM/Ensenada for access to their Origin-2000 computer.

#### REFERENCES

- Aguilar, L. A., & Hoogerwerf, R. 1999, IAU Symp. 186, p.47-51.  
 de Bruijne, J. H. 1999, MNRAS 306, p.381.  
 de Zeeuw, P. T., Hoogerwerf, R., de Bruijne, J. H., Brown, A. G., & Blaauw, A. 1999, AJ 117, p.354-399  
 ESA, 1997, ESA SP-1200, The *Hipparcos* and *Tycho* Catalogues. ESA Publications.  
 Hoogerwerf, R., & Aguilar, L. A. 1999, MNRAS 306, p.394-406.  
 Perryman, M. A., et al., 1998, A&A 331, p.81-120