

## KINEMATICS AROUND THE NON-THERMAL SUPERBUBBLE IN IC 10

A. Bullejos<sup>1</sup> and M. Rosado<sup>1</sup>

**We study the kinematics around a non-thermal superbubble found by Yang & Skillman. Considering the H II and [S II] profiles, we find that between 3 and 6 supernovae are required to form this superbubble.**

IC 10 may well be the nearest example of a blue compact dwarf galaxy (BCD). Among the Local Group star-forming dwarf galaxies, IC 10 is conspicuous for having the highest surface density of WR stars ( $11.2 \text{ kpc}^{-2}$ , Richer et al. 2000). The presence of so many WR stars and the high H $\alpha$  luminosity emphasize that IC10 is currently undergoing a burst of star formation that began at least 10 Myr ago. Observations of 21-cm emission from neutral hydrogen reveal that IC 10 consists of an inner disk embedded in an extended, complex, counter-rotating envelope (Shostak & Skillman 1989). This leads Wilcots & Miller (1998) to conclude that IC 10 is still in its formative stage. These H I observations make clear the youth of the current star formation episode for there is a notable lack of interstellar medium structures attributable to supernovae (Wilcots & Miller 1998). The galaxy size in the light of H $\alpha$  is  $5 \times 3$  arcmin while in H I it is  $18 \times 15$  arcmin. All optical studies of IC 10 are hampered by the large foreground reddening due to its position near the plane of our galaxy ( $l = 119^\circ$ ,  $b = -3^\circ$ ).

Our observations of the non-thermal superbubble, discovered by Yang & Skillman (1993), were obtained in the light of H $\alpha$  and [S II] with the scanning Fabry-Perot Interferometer, PUMA. We suppose that the energy in the ionized interstellar medium comes from the energy injected by stellar winds and supernovae. Considering both the H $\alpha$  and [S II] profiles, we find a typical expansion velocity of 50 to 70  $\text{km s}^{-1}$ . Using a distance of 661 kpc (Sakai et al. 1999) and a diameter of  $41.5''$  inferred from Yang & Skillman (1993), we calculate a radius of 66 pc for the superbubble. Adopting a density of  $1.1 \text{ atoms cm}^{-3}$  (Yang & Skillman 1993), the mass swept out by the superbubble is  $4.87 \times 10^{37} \text{ g}$  ( $2.45 \times 10^4$  solar masses).

<sup>1</sup>Instituto de Astronomía, UNAM, Apdo Postal 70-264, Ciudad Universitaria, México D.F., México (almudena,margarit@astroscu.unam.mx).




Fig. 1. A close-up of the region of the non-thermal superbubble in H $\alpha$  with the Yang & Skillman, (1993) iso-contours of the 49-cm flux superposed. This image was formed by summing up the velocity channels containing emission from IC 10. The heliocentric velocity of this galaxy is  $-344 \text{ km s}^{-1}$  (de Vaucouleurs et al. 1991).

With these values, the kinetic energy obtained is  $6.09 \times 10^{50}$  to  $11.9 \times 10^{50}$  erg. Supposing that 20% of a supernova's thermal energy is converted into kinetic energy and that each of them injects  $10^{51}$  erg, then from 3 to 6 supernova are required to form this superbubble.

### REFERENCES

- de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H. G., et al. 1991, in *The Third Reference Catalog of Bright Galaxies* (New York: Springer)
- Richer, M. G. et al. 2001, *A&A*, 370, 34
- Sakai, S., Madore, B.F., & Freedman, W. L. 1999, *ApJ*, 511, 671
- Shostak, G. S., & Skillman, E. D. 1989, *A&A*, 214, 33
- Wilcots, E. M., Miller, B. W. 1998, *AJ*, 116, 2362
- Yang, H., & Skillman, E. D. 1993, *AJ*, 106, 1448