

NEW LIGHT ON THE SEARCH FOR LOW-METALLICITY GALAXIES

G. Denicoló,¹ R. Terlevich,^{1,2} and E. Terlevich^{3,4}

In many cases the oxygen abundance can only be estimated using empirical methods such as R_{23} (Pagel et al. 1979), S_{23} (Vílchez & Esteban 1996) or S_{234} (Oey & Shields 2000). But in face of the weak points and limitations of each R_{23} and $S_{23(4)}$ methods, we understand that a more precise calibration with another method involving easily measurable lines, for a larger redshift range and with a monotonic low-dispersion relation, would be highly desirable.

In this work we have used the best available data to explore the usefulness of the N_2 abundance calibrator, where $N_2 = [\text{N II}] \lambda 6584/\text{H}\alpha$ (Storchi-Bergmann et al. 1994). The metal-rich galaxy sample includes data from Díaz et al. (2000a,b), Terlevich et al. (1991).

We have used the N_2 estimator to select low metallicity candidates from redshift survey quality data. We have found six metal-poor galaxies ($7.6 < 12 + \log \text{O/H} < 8.0$) selected from the Durham/UKST Galaxy Redshift Survey (Ratcliffe et al. 1998). We find that metal-poor galaxies represent only a small fraction (0.2%) of the total observed emission line galaxies. Our next approach is to use the N_2 estimator in a large redshift survey of galaxies such as 2dF aiming to significantly increase the present sample of metal-poor galaxies and perhaps more importantly to study the cosmological abundance evolution. In an attempt to quantify the dispersion in the N_2 relation, we have studied the connection between the scatter of the N_2 parameter with the equivalent width of $\text{H}\beta$. We find a slight tendency for galaxies with positive residuals to have the largest $\text{EW}(\text{H}\beta)$. We do not report any other trend of the residuals with $\text{EW}(\text{H}\alpha)$ or with $\log([\text{O II}]/[\text{O III}])$.

¹Institute of Astronomy, Madingley Road, Cambridge, CB3 0HA, UK.

²Visiting Professor, INAOE, Puebla, Mexico.

³Instituto Nacional de Astrofísica, Óptica y Electrónica, Tonantzintla, Puebla, Mexico.

⁴Visiting Fellow, IoA, Cambridge.




Fig. 1. The N_2 metallicity calibrator is defined as $N_2 = [\text{N II}] \lambda 6584/\text{H}\alpha$. The linear fit to the data, taking into account the dispersion in the distribution, gives $12 + \log \text{O/H} = 9.12 + 0.73 \log(N_2)$, with a correlation coefficient of 0.85.

This work is discussed more fully in Denicoló, Terlevich, & Terlevich (2002).

REFERENCES

- Díaz, A. I., & Pérez-Montero, E. 2000, MNRAS, 312, 130
- Díaz, A. I., Castellanos, M., Terlevich, E., & Luisa García-Vargas, M. 2000, MNRAS, 318, 462
- Oey, M. S., & Shields, J. C. 2000, ApJ, 539, 687
- Pagel, B. E. J., Edmunds, M. G., Blackwell, D. E., Chun, M. S., & Smith, G. 1979, MNRAS, 189, 95
- Ratcliffe, A., et al. 1998, MNRAS, 300, 417
- Storchi-Bergmann, T., Calzetti, D., & Kinney, A. L. 1994, ApJ, 429, 572
- Terlevich, R. J., Melnick, J., Masegosa, J., Moles, M., Copetti, M. V. F. 1991, A&A, 91, 285
- Denicoló, G., Terlevich, R. J., & Terlevich, E. 2002, MNRAS, 330, 69
- Vílchez, J. M., & Esteban, C. 1996, MNRAS, 280, 720