

THE MASS PROFILE OF NGC 1399 DETERMINED FROM GLOBULAR CLUSTER DYNAMICS

B. Dirsch¹, T. Richtler¹, D. Geisler¹, K. Gebhardt², L. Infante^{3,4}, D. Minniti⁴, M. Hilker⁵, M. Rejkuba⁴, E.K. Grebel⁶, S. Larsen⁷ and M.V. Alonso⁸

Spectra of hundreds of globular clusters around NGC 1399 have been acquired at the VLT with FOR S2 / MXU. 470 cluster radial velocities have been obtained, representing the largest velocity sample for any galaxy collected so far. This sample is large enough to allow a fully self-consistent dynamical modeling. We present our sample and give preliminary results.

Globular clusters have a high potential to study the mass profile of early-type galaxies out to large radii using them as dynamical tracers. NGC 1399, the central galaxy of the Fornax cluster, is a particularly promising target for such an investigation due to its rich cluster system (≈ 6000 , e.g. Ostrov et al. 1998). We obtained 470 cluster radial velocities with a mean uncertainty of 55 km s^{-1} . The uncertainty has been derived by comparing independent velocity measurements for a subset of clusters on different masks. They cover the radial range from $2'$ to $8'$ corresponding to 11 kpc to 44 kpc. The velocities have been obtained with cross-correlation techniques as well as with direct line measurements. Additionally, photometry in the Washington system has been acquired that allows straightforward identification of a red, metal-rich and a blue, metal-poor cluster population due to the strong bimodality of the color distribution.

A full dynamical analysis of the data is underway and will be published soon.

We observed a radially constant velocity dispersion ($304 \pm 11 \text{ km s}^{-1}$ for the whole sample) which is in contrast to former studies that used much smaller velocity samples (Kissler-Patig et al. 1999). Moreover, we found that the metal-rich and the metal-

poor subpopulations have a significantly different *projected* velocity dispersions of $297 \pm 16 \text{ km s}^{-1}$ and $355 \pm 22 \text{ km s}^{-1}$, respectively.

In our analysis we assumed a spherically symmetric three dimensional density distribution of globular clusters. With a power law of the form $\rho \propto r^\alpha$ we find for red and blue subpopulations $\alpha = -2.5 \pm 0.1$ and $\alpha = -1.7 \pm 0.1$, respectively (assuming spherical symmetry). Under the additional assumption of no-rotation the difference in the measured velocity dispersions is in accordance with the density profile difference assuming isotropy (this can be seen from the spherical Jeans equation, e.g. Binney & Tremaine 1997). The radially constant velocity dispersion implies an r^{-2} mass distribution, representing an isothermal sphere. This corresponds to a constant circular velocity of $460 \pm 25 \text{ km s}^{-1}$. For such a mass distribution the mass-to-light ratio measured in the B band rises from $20 M_\odot/L_\odot$ at 11 kpc to $40 M_\odot/L_\odot$ at 44 kpc. The total mass within 40 kpc is in good agreement with the mass determinations based on ROSAT data (Jones et al. 1997)

BD acknowledges financial support by the Alexander von Humboldt Society.

REFERENCES

- Binney J. & Tremaine S. 1997, Galactic Dynamics, Princeton Series in Astrophysics
Ostrov P.G., Forte J.C., & Geisler D. 1998, AJ, 116, 2854
Jones C., Stern C., & Forman W. 1997, ApJ, 482, 143
Kissler-Patig, M., Grillmair C.J., Meylan G., et al. 1999, AJ, 117, 1206

¹Universidad de Concepción, Casilla 160-C, Concepción, Chile

²University of Texas, 26th and Speedway, Austin, TX 78712, USA

³Universidad Católica, Santiago, Chile

⁴ESO, Garching, Alemania

⁵Sternwarte der Universität Bonn, Bonn, Alemania

⁶MPIA, Heidelberg, Alemania

⁷Lick Observatory, UC Santa Cruz, USA

⁸Observatorio Astronómico de Córdoba, UNC, Argentina