

MODELING THE BUILDING BLOCKS OF WATER MASERS IN CEPHEUS A

L. Uscanga,¹ J. Cantó,¹ S. Curiel,¹ J. M. Torrelles,² and L. F. Rodríguez³

Recent VLBA observations of water maser emission from the star-forming region Cepheus A have shown that this emission actually comes from a set of coherent features or “building blocks”.

These building blocks are small linear structures of water maser spots that conform a larger configuration with a well-defined linear/arcuate shape. In small angular regions, the VLBA observations (Torrelles et al. 2001) show that:

1. The water maser spots form a well-defined linear chain in the plane of sky.
2. A single spectral line with a width of 0.5 to 1 km s⁻¹ in the flux density vs. velocity graph.
3. A coherent structure and a well-defined velocity gradient in the position-velocity diagrams.

A conspicuous example of these results is the R5-a block containing the strongest maser spot of the entire region (~ 200 Jy beam⁻¹, see Figure 1).

This building block is modeled as a thin disk of radius r , and thickness h , of material flowing in the direction θ with a linear velocity gradient a . The observer is in the disk plane.

The following equation was used to express the intensity from each point of the disk at a given velocity:

$$I(v, y, z) = I_0 \exp \left\{ \int \kappa_0 e^{-(v-v_r)^2/\Delta v^2} dx \right\}, \quad (1)$$

where the term in brackets is the optical depth and v_r is the radial velocity of the flow at the position (x, y) . I_0 , κ_0 , and Δv are supposed homogeneous throughout the disk. Once the parameters are specified, we compute the flux density, $F(v)$, and the coordinates of the emission centroid at velocity v , $y_c(v)$ and $z_c(v)$. The results are compared with the observed maser spots of the R5-a block (Figure 2).

REFERENCES

Torrelles, J. M., et al. 2001, ApJ, 560, 853

¹Instituto de Astronomía, UNAM, Apdo. Postal 70-264, 04510 México, D.F., México (lucero@astroscu.unam.mx).

²Institut d’Estudis Espacials de Catalunya (IEEC/CSIC), Spain.

³Instituto de Astronomía, UNAM, Morelia, México.




Fig. 1. (a) R5 water maser offset positions on 1996 February 11. The dash line shows the least-squares fitted circle. (b) Close-up of the R5-a block. The sizes of the circles are proportional to the maser spot intensity.




Fig. 2. (a) Emission centroid positions on the plane of the sky, (b) flux density as a function of velocity, and (c) position-velocity diagram: solid line—model, dots—observations. The observed positions are measured along an axis with position angle $\simeq -63^\circ$. The zero position corresponds to the geometrical center of the R5-a block. Model parameters: $I_0 = 2.25$ Jy beam⁻¹ AU⁻², $\kappa_0 = 2.24$ AU⁻¹, $\Delta v = 0.67$ km s⁻¹, $v_0 = 2.85$ km s⁻¹, $r = 1.5$ AU, $h = 0.15$ AU, $a = 0.29$ km s⁻¹ AU⁻¹, $\theta = 9^\circ$.