

STATISTICS AND MULTIWAVELENGTH SYNTHESIS MODELS: TOWARDS A NEW GENERATION OF SYNTHESIS MODELS

M. Cerviño^{1,2}

In this contribution I present my current work on a new generation of evolutionary synthesis models that compute the multiwavelength energy distribution (from gamma rays to radio) as well as the associated dispersion for young stellar systems. I will also show some statistical effects that may appear in the analysis of surveys, such as bimodal or multimodal distributions and bias when color indices computed by the codes are compared with observations. Such a new generation of synthesis models may be useful for the analysis of the data expected from the GTC.

1. INTRODUCTION

Taking into account the discreteness of the stellar population, the predictions of any model that relies on an initial mass function (IMF) are exact only *under the assumption of an infinite number of stars*. Otherwise, they only give a *mean value* of a probability distribution. The relevance of such fluctuations in the results of synthesis models is obvious in the case of massive stars and young clusters, *but they also affect the models of older clusters dominated by the emission of low-mass stars* since small variations in the initial mass/number of stars in a given mass range can produce different numbers of, for example, AGB stars at a given age, which in turn produce large variations in the resulting observable.

So for the comparison of models with observational data it is necessary to obtain not only the mean value of the observable but also, at least, the corresponding dispersion of the computed observable due to the discreteness of the stellar population, and, ideally, the underlying probability density distributions. The dispersion due to a finite number of stars in real systems can be evaluated theoretically, as has been shown in Cerviño et al. (2002).

2. SOME EXAMPLES: MULTIWAVELENGTH EMISSION AND COLOR-COLOR DIAGRAMS

In Figure 1 I show the 90% confidence level for the multiwavelength spectrum for a 5.5 Myr old

burst obtained analytically from our code. Examples of statistical effects of a large number of Monte Carlo simulations and comparisons with the dispersion obtained theoretically for different observables can be found in Cerviño et al. (2000, 2001, 2002).

In Figure 2 I show the $B - V$ vs. $V - K$ colors for a 10 Myr old star forming region. Note that *it is assumed that colors are independent on the number of stars in the cluster*, so I have tested this hypothesis with 10^3 Monte Carlo simulation of clusters (triangles) with 1000, 100, 10, and 1 stars in the mass range 2–120 M_{\odot} following a Salpeter IMF slope. The theoretical result is shown with the \odot symbol. The figure shows two important effects: a) There are bimodal (multimodal) distributions of the colors; b) the mean value obtained by infinite populated IMF (analytical result) has a bias if it is used for the analysis clusters with a small number of stars. As reference, the mean mass of a cluster with 10^3 stars in the given mass limits is $\approx 6 \times 10^3 M_{\odot}$, it means $2 \times 10^4 M_{\odot}$ if the value of lower mass limit of the IMF is $0.08 M_{\odot}$. Both, bimodality, and bias are a natural effect of statistics when only a few stars dominate the observable. Note that it is dependent on the observable: *Example 1:* A cluster with 10^5 stars in the mass range 2–120 M_{\odot} with a Salpeter IMF slope have about only 10 Wolf-Rayet stars! *Example 2:* The effect on the colors of globular clusters (dominated by a few AGB stars) can be relevant (see Bruzual 2001 and references therein for details). Additionally, these bimodality effects produce a bias when ratios and logarithm quantities computed by synthesis codes are compared with observed data (but such a bias is not present for quantities such as luminosities even for “clusters” with only 1 star). This subject is addressed in Cerviño & Valls-Gabaud (submitted).

3. DISCUSSION

Statistical effects may be especially relevant: i) In the analysis of individual systems, in particular for the galaxy building blocks if they are small clusters; ii) in the analysis of the integrated spectra obtained with high resolution instruments when not all the region is covered by the slit; iii) in chemical evolutionary models (Cerviño & Mollá,

¹LAEFF, INTA, Apdo. 50727, E-28080 Madrid, Spain.

²Instituto de Astrofísica de Andalucía, CSIC, Camino Bajo de Huétor 24, E-18080 Granada, Spain.




Fig. 1. Analytical 90% confidence level for the multiwavelength spectrum for a 5.5 Myr old burst.




Fig. 2. Monte Carlo simulations of the $B - V$ vs. $V - K$ colors of 10 Myr old stellar clusters with 1000, 100, 10, and 1 stars in the mass range 2–120 M_{\odot} following a Salpeter IMF slope (triangles). The theoretical result is showed with the \odot symbol.

submitted), in so far as they are dependent on the yields produced from supernovae, which are intrinsically “rare” events; iv) in the analysis of surveys, in the special case of galaxies with emission

lines produced by massive stars that are intrinsically a small fraction of the total number of stars but dominate the emission. In all these cases, the use of multiwavelength observations and self-consistent synthesis models that include such statistical effects and compute the correlation between different wavelengths will be an asset for safely constraining the physical properties of stellar systems (from small star forming regions to high redshift galaxies). The current model outputs can be found in <http://www.laef.esa.es/mcs/SED/>.

REFERENCES

- Bruzual, G. 2001 in IAU Symp. 207, Extragalactic Star Clusters, ed. E. K. Grebel, D. Geisler, & D. Minniti (San Francisco: ASP), in press
- Cerviño, M., Luridiana, V., & Castander, F. J. 2000, *A&A*, 360, L5
- Cerviño, M., Gómez-Flechoso, M. A., Castander, F. J., Schaerer, D., Mollá, M., Knödseder, J., & Luridiana, V. 2001, *A&A*, 376, 422
- Cerviño, M., Valls-Gabaud, D., Luridiana, V., & Mas-Hesse, J. M. 2002, *A&A*, 381, 51