

## THE CONTRIBUTION OF X-RAY SELECTED SOURCES TO THE SUB-MILLIMETER BACKGROUND

P. Severgnini,<sup>1</sup> R. Maiolino,<sup>2</sup> M. Brusa,<sup>3</sup> A. Cimatti,<sup>2</sup> A. Comastri,<sup>4</sup> F. Fiore<sup>5</sup> and C. Vignali<sup>6</sup>

**The connection between the sub-millimeter (sub-mm) and hard X-ray background (XRB) has been studied by comparing data at 2–10 keV and at 850  $\mu\text{m}$  for a sample of 54 sources. At the limiting fluxes reached in our sample most of the background ( $\sim 70\%$ ) in the two bands has been resolved into discrete sources. We find that sources making most of the X-ray background do not contribute significantly to the sub-mm background.**

**The Sample:** The objects included in our program are two sub-samples of hard X-ray sources detected by Beppo-SAX in the HELLAS survey (Fiore et al. 2001) and by *Chandra* in the deep hard X-ray survey of the Hawaii Field SSA13 (Mushotzky et al. 2000). The X-ray and sub-mm observed fluxes and redshifts for the selected sources are reported in Table 1. For these objects our own SCUBA observations have been carried out. The SCUBA observations and data reduction are described in details in Severgnini et al. (2000). None of the sources has been detected at 850  $\mu\text{m}$ .

TABLE 1

Source	$F_{2-10\text{keV}}$ [ $10^{-13}$ cgs]	$F_{850}^a$ [mJy]	z
Beppo-SAX 0045-2515	3.5	<2.6	0.111
1054+5725	2.7	<3.4	0.205
1117+4018	1.3	<7	1.274
2302+0856	3.3	<2.4	0.135
<i>Chandra</i> 1312+4239	0.27	<3.2	1.048
1312+4240	0.042	<3	...
1312+4241	0.23	<4.4	1.32
1312+4242	0.049	<2.8	2.415

<sup>a</sup> Upper limits are at  $2\sigma$ .

We have combined our two sub-samples with other 46 sources obtained by cross-correlating sub-mm and X-ray data available from the literature

<sup>1</sup>INAF - Osservatorio Astronomico di Brera, Via Brera 28, I-20121 Milano, Italy (paola@brera.mi.astro.it).

<sup>2</sup>INAF - Osservatorio Astronomico di Arcetri, Italy

<sup>3</sup>Dipartimento di Astronomia Università di Bologna, Italy.

<sup>4</sup>INAF - Osservatorio Astronomico di Bologna, Italy.

<sup>5</sup>INAF - Osservatorio Astronomico di Roma, Italy.

<sup>6</sup>Department of Astronomy and Astrophysics, Pennsylvania State University, USA.

(Mushotzky et al. 2000, Barger et al. 1999, 2001a; Fabian et al. 2000; Hornschmeier et al. 2000; Bautz et al. 2000). The whole sample here considered comprises thus 54 objects.

**Results:** For each source we have computed the sub-mm-to-X-ray index  $\alpha_{SX}$  ( $F_\nu \propto \nu^{-\alpha}$ ), using the rest-frame flux densities at 850  $\mu\text{m}$  and 5 keV (or upper limits if the source was not detected). By comparing the  $\alpha_{SX}$  as a function of redshift for our whole sample with that of various classes of templates (Type 1 unobscured AGN, Type 2 obscured AGN and starburst galaxies) and with that of the Cosmic Background, constraints have been given on the contribution of the hard X-ray population to the sub-mm background. In particular, we find that, under conservative assumptions, the contribution to the sub-mm background from 2–10 keV sources brighter than  $\sim 10^{-15}$  erg s<sup>-1</sup>cm<sup>-2</sup> (which resolve at least 75% of the background in this band), is about 7%. Recent *Chandra* observations (Brandt et al. 2001; Barger et al. 2001b) have extended our results to the X-ray sources making about 90% of the hard XRB, and find that their contribution to the sub-mm background is about 15%. This value, derived with better statistics and obtained with a different approach, is in agreement with our results. Any significant contribution to the sub-mm background is limited to fainter hard X-ray sources which might contribute at most for 10% of the 2–10 keV background. Finally, our estimate is also in good agreement with the recent models predicting the contribution of X-ray sources to the sub-mm background (see Brusa et al. 2001 and references therein).

## REFERENCES

- Brusa M., et al. 2001 (astro-ph/0106014)
- Barger et al. 1999, ApJ 518, L5
- Barger et al. 2001a, AJ 121, 662
- Barger et al. 2001b, ApJ 560, L23
- Bautz et al. 2000, AJ 543, L119
- Brandt et al. 2001, AJ 122, 2810
- Fabian et al. 2000, MNRAS 315, L8
- Fiore et al. 2001, MNRAS 327, 771
- Hornschmeier et al. 2000, ApJ 541, 49
- Mushotzky et al. 2000, Nature 404, 459
- Severgnini et al. 2000, A&A 360, 457