

ABSTRACTS OF CONTRIBUTED PAPERS

THE 2003 OUTBURST OF V4641 SGR (=SAX 1819.3-2525) Charles Bailyn¹

V4641 Sgr (=SAX 1819.3-2525) is a transient black hole binary system, with a B-type secondary star. In August 2003 a new outburst of this source began. Violent variability (factors of 2 on timescales of 10 seconds or less) was seen in all wavelength regimes (radio, infrared, optical and X-ray). I will present data from our RXTE observations and OIR data from the SMARTS 1.3m telescope at CTIO (ATEL 171, 172). Observations continue as of this writing (August 8, 2003).

¹ Yale University, USA

THE NATURE OF ULTRALUMINOUS SOURCES: X-RAY AND OPTICAL CLUES Manfred W. Pakull¹

It is proposed to review recent XMM and optical (CFHT/ SUBARU) observations of ultraluminous X-ray sources in nearby galaxies, including the apparently brightest ULXs which are located in the M81 group dwarf galaxies Holmberg II and IX. We show that their 0.2-10 keV spectra can be characterised by simple power-law models and by negligible contribution of any (disk) thermal component, if the low-metallicity of the interstellar absorber is taken into account. Optical observations have revealed the presence of large ionized nebulae around many ULX which provide important informations on intrinsic luminosities, and on the formation and evolution of these sources. Finally, the first detection of a stellar emission line in the spectrum of an optical counterpart will be presented, and the prospects to determine the mass of an ULX will be discussed.

¹ Observatoire de Strasbourg, France

JET INDUCED NUCLEOSYNTHESIS IN MISALIGNED MICROQUASARS Yousaf Butt¹

The jet axes and the orbital planes of microquasar systems are usually assumed to be approximately perpendicular, even though this is not currently an observational requirement. On the contrary, in one of the few systems where the relative orientations are well-constrained, V4641Sgr, the jet axis is known to lie not more than 36 degrees from the binary plane. Such a jet, lying close to the binary plane, and traveling at a significant fraction of the speed of light may periodically impact the secondary star initiating nuclear reactions on its surface. The integrated yield of such nuclear reactions over the age of the binary system (less the radiative mass loss) will detectably alter the elemental abundances of the companion star. This scenario may explain the anomalously high Li enhancements (roughly 20-200 times the sun's photospheric value; or, equivalently, 0.1-1 times the average solar system value) seen in the companions of some black-hole X-ray binary systems. Gamma-ray line signatures of the proposed process could include the 2.22 MeV neutron capture line as well as the 0.478 MeV ⁷Li* de-excitation line, both of which may be discernable with the INTEGRAL satellite if produced in an optically thin region during a large, close-by outburst. For very energetic jets, a relatively narrow neutral pion gamma-decay signature at 67.5 MeV could also be measurable with the GLAST satellite. We argue that about 10-20% of all microquasar systems ought to be sufficiently misaligned as to be undergoing the proposed jet-secondary impacts.

¹ CfA, USA

THEORETICAL MODELS OF X-RAY BINARIES IN NEARBY GALAXIES

Vicky Kalogera¹

In recent years Chandra has revolutionized the study of X-ray binaries with detection of significant samples in a wide range of galaxies in the nearby Universe. Most often these populations are characterized by X-ray luminosity functions which exhibit systematic behaviors correlated with galaxian properties. We have recently developed theoretical models of X-ray binary formation and evolution that allow for the first time the direct comparison with X-ray observations. Our first results related to star-forming galaxies are very encouraging and clearly show that they can confirm population properties (for example ages) derived based on observations in other wavelengths and they allow to derive physical information about X-ray binaries in diverse environments.

¹ Northwestern University, USA

GALACTIC POPULATIONS OF ULTRACOMPACT BINARIES

Krzysztof Belczyński¹

Using the StarTrack population synthesis code we analyze the formation and evolution of ultracompact binaries (UCBs). We study UCBs formed only in the Galactic field and focus on systems with a neutron star or a black hole accretor. We predict the relative numbers of different types of UCBs populating our Galaxy. We find an excellent agreement with observed sample of Galactic UCBs. Our results demonstrate that regular binary star evolution may lead to formation of all UCBs in the Galactic field. No dynamical processes of the dense stellar environments (e.g. globular clusters) are necessary to explain formation of the field UCBs. We predict that with a discovery of several more X-ray active UCBs hosting neutron stars, we should find the system with a black hole accretor.

¹ Northwestern University, USA

THE EXTRAORDINARY LUMINOUS BINARY STAR, QU CAR

Louise Hartley¹

In recent UV and optical studies the interacting binary star, QU Car, has been shown to be extremely bright and highly ionised – a unique system amongst CVs. Assigned to the nova-like subclass of CVs it is, however, an order of magnitude more luminous than is typical. Furthermore, it displays spectral characteristics that would suggest a close similarity to the supersoft X-ray sources, although it is several orders of magnitude too faint in the soft X-ray range to be a supersoft. It has also shown to be the first known CV with a carbon-enriched companion. Here we present spectroscopic observations of QU Car, from the far ultraviolet through to the optical range. We show how we have built up a picture of QU Car, including its distance; its extremely high mass accretion rate; and its unusual ionisation and abundance patterns. We then discuss QU Car's relation to other luminous interacting binaries, including the supersoft sources and V Sge systems, and highlight the rewards that are to be gained from understanding the evolutionary path of this system.

¹ Monash University, USA

TYPE IA SUPERNOVAE AND THEIR IMPLICATIONS FOR COSMOLOGY

Mario Livio¹

Models for Type Ia Supernovae (SNe Ia) are reviewed. It is shown that there are strong reasons to believe that most SNe Ia represent thermonuclear disruptions of C-O white dwarfs, when these white dwarfs reach the Chandrasekhar limit and ignite carbon at their centers.

Different progenitor scenarios are reviewed critically and the strengths and weaknesses of each scenario are presented in detail. It is argued that theoretical considerations currently favor single-degenerate models, in which the white dwarf accretes from a subgiant or giant observed sample. The recent observations of SN 2002ic and their implications are reviewed. The relation of the different models to the use of SNe Ia for the determination of cosmological parameters is discussed. It is shown that while the observed diversity of SNe Ia may argue for the existence of different progenitor classes, this does not affect the interpretation of an accelerating expansion of the universe.

Crucial observations tests of the conclusions are suggested.

¹ Space Telescope Science Institute, USA

RELATIVISTIC MEASUREMENTS OF
PULSAR-WHITE DWARF BINARIES: NEW
RESULTS FROM ARECIBO

D. Nice¹

High precision timing observations of binary radio pulsars in nearly circular orbits with white dwarf companions have lead to new measurements and constraints on orbital geometries and component star masses. These constraints come primarily from measurements of relativistic phenomena, including precession, Shapiro delay, and orbital decay due to emission of gravitational radiation. Masses of white dwarf secondaries in these systems show good agreement with expectations from the theoretical orbital period-core mass relation. Masses of neutron star primaries in these systems show a trend of values greater than the canonical 1.35 solar masses, presumably due to mass accretion during the spin-up of the pulsars.

¹ Princeton University, USA

MULTIWAVELENGTH OBSERVATIONS OF RX
J0806.3+1527: THE SHORTEST ORBITAL
PERIOD BIN

G. Israel¹

I will report on the results obtained from observations carried out with XMM-Newton, the Very Large telescope (VLT) and the Telescopio Nazionale

Galileo (TNG) covering the X-ray, EUV, optical and IR bands. These observations allowed us to infer new important results related to the stability of the absolute phase shift between Optical/IR and X-ray emission, the X-ray spectrum as a function of the orbital phase, the period derivative, the polarisation, and the EUV emission. These results will be discussed on the light of the theoretical models proposed so far for RXJ0806.3+1527 and its twin source RXJ1914.4+2456

¹ INAF - OAR, Italy

X-RAY OBSERVATIONS OF NOVA V4743
SAGITTARIUS IN OUTBURST

J. Krautter,¹ J.-U. Ness,¹ S. Starrfield,¹
V. Burwitz,¹ J. J. Drake,¹ and M. Orio¹

The results of Chandra and XMM-Newton X-ray observations of V4743 Sgr (Nova Sgr 2002 no.2) carried out on several epochs after outburst will be presented here. Six months after outburst the nova had evolved into a Super-Soft X-ray source exhibiting a continuous spectrum with deep absorption features. The X-ray light curve showed large amplitude oscillations with a period of 1325 sec followed by a deep decline after about 13 ksec of observations. The count rate dropped from 40 cts/sec to practically zero with 6 ksec. The spectral hardness changed in correlation with the oscillations and became significantly softer during the decline. Variations of this strength of the X-ray flux had never been observed before in any nova in outburst. Possible mechanisms for the variations will be discussed.

¹ Landessternwarte Heidelberg, Germany

LIST OF ABSTRACTS

THE 2003 OUTBURST OF V4641 SGR (=SAX 1819.3-2525) <i>C. Bailyn</i>	273
THE NATURE OF ULTRALUMINOUS SOURCES: X-RAY AND OPTICAL CLUES <i>M. W. Pakull</i>	273
JET INDUCED NUCLEOSYNTHESIS IN MIS- ALIGNED MICROQUASARS <i>Y. Butt</i>	273
THEORETICAL MODELS OF X-RAY BINARIES IN NEARBY GALAXIES <i>V. Kalogera</i>	274
THE EXTRAORDINARY LUMINOUS BINARY STAR, QU CAR <i>L. Hartley</i>	274
GALACTIC POPULATIONS OF ULTRACOM- PAC BINARIES <i>K. Belczyński</i>	274

TYPE IA SUPERNOVAE AND THEIR IMPLICA- TIONS FOR COSMOLOGY <i>M. Livio</i>	274
RELATIVISTIC MEASUREMENTS OF PULSAR- WHITE DWARF BINARIES: NEW RESULTS FROM ARECIBO <i>D. Nice</i>	275
MULTIWAVELENGTH OBSERVATIONS OF RX J0806.3+1527: THE SHORTEST ORBITAL PE- RIOD BIN <i>G. Israel</i>	275
X-RAY OBSERVATIONS OF NOVA V4743 SAGITTARIUS IN OUTBURST <i>J. Krautter, J.-U. Ness, S. Starrfield, V. Burwitz, J. J. Drake, & M. Orio</i>	275