

OLD MAIN-SEQUENCE TURNOFF PHOTOMETRY IN THE SMC

N. E. D. Noël,¹ C. Gallart,¹ E. Costa,² and R. A. Méndez²

RESUMEN

Se presenta un conjunto de diagramas color-magnitud (DCM) correspondientes a doce campos localizados en la Pequeña Nube de Magallanes (SMC). Estos DCM poseen una profundidad sin precedentes, llegando hasta los puntos de giro más viejos de la secuencia principal y cubriendo un amplio rango de distancias galactocéntricas, hasta $\sim 4^\circ$ desde el centro de la SMC y con diferentes ángulos de posición. Se expone un análisis del contenido estelar en estos campos de la SMC a través de la comparación con isocronas teóricas. Nuestro estudio confirma la existencia de fuertes gradientes de población y variaciones espaciales en el contenido estelar de la SMC. Ninguno de los campos mostrados está dominado por una población estelar exclusivamente vieja. Esto prueba que a $\sim 4^\circ$ del centro de la SMC no hemos alcanzado un halo estelar viejo similar al de la Vía Láctea.

ABSTRACT

We present ground-based *B* and *R*-band color-magnitude diagrams (CMDs) of unprecedented depth for twelve fields in the Small Magellanic Cloud (SMC). They reach the oldest main-sequence turnoffs and cover a wide range of galactocentric distances up to $\sim 4^\circ$ from the SMC center, and are located at different position angles. A picture of the stellar content in our SMC fields is presented, through the comparison with theoretical isochrones. Our study confirms the existence of strong population gradients and spatial variation in the SMC stellar content. None of the SMC fields presented here are dominated by old stellar populations which proves that at $\sim 4^\circ$ from the SMC center we do not reach an old stellar halo similar to that of the Milky Way.

Key Words: GALAXIES: EVOLUTION — GALAXIES: INDIVIDUAL (SMC) — GALAXIES: STELLAR CONTENT — LOCAL GROUP

1. INTRODUCTION

Besides the intrinsic interest in studying the SMC stellar populations, which is key to understand our local environment, this galaxy has been largely neglected in favour of its closest neighbour, the LMC. There is a disagreement in the published SFHs of the SMC since some authors (e.g. Harris & Zaritsky 2004) argue that there was a quiescent period at intermediate ages in which the galaxy formed few or no stars, and others (e.g. Dolphin et al. 2001) do not find a gap in the star formation rate of the SMC. It is important to stress that the SMC regions analysed by these authors are different. In an attempt to address this conflict, here we present a discussion based on deep CMDs with theoretical isochrones overlapped.

2. SMC STELLAR CONTENT

The interpretation of composite stellar population CMDs strongly relies on the stellar evolution models adopted (see Gallart et al. 2005). For our

purpose we used the Teramo stellar evolution models (Pietrinferni et al. 2004) as shown in Figure 1 (Eastern SMC fields in the upper panel and Western SMC fields in the lower panel) and in Figure 2 (Southern SMC fields), using metallicities suitable for the SMC stellar populations.

Our analysis shows the presence of spatial variations in the stellar content as a function of the position angle and strong gradients in the stellar population as a function of the galactocentric distance. The fact that the SMC CMDs analyzed here do not show a blue horizontal branch and that none of them is dominated by a completely old population indicate that, at $\sim 4^\circ$ from the SMC center, we do not reach an old halo similar to that of the Milky Way (Noël et al. 2006).

REFERENCES

- Dolphin, A. E., Walker, A. R., Hodge, P. W., Mateo, M., et al. 2001, *ApJ*, 562, 303
Gallart, C., Zoccali, M., & Aparicio, A. 2005, *ARA&A*, 43, 10
Harris, J., & Zaritsky, D. 2004, *AJ*, 127, 1531
Noël, N. E. D., Gallart, C., Costa, E., & Méndez, R. A. 2006, *AJ*, submitted
Pietrinferni, A., Cassisi, S., Salaris, M., & Castelli, F. 2004, *ApJ*, 612, 168

¹Instituto de Astrofísica de Canarias, 38200, La Laguna, Tenerife, Spain (noelia@iac.es).

²Universidad de Chile, Casilla 36-D, Las Condes, Santiago, Chile.




Fig. 1. Left panel shows the CMDs of the Eastern SMC fields, located in a range of $\sim 1.4^\circ$ to $\sim 1.7^\circ$ from the center. Canonical isochrones of 6, 9, and 13 Gyr with $Z=0.001$, isochrones with overshooting of 1, 3, and 4 Gyr with $Z=0.002$, and of 0.1 Gyr and 0.03 Gyr with $Z=0.004$ were overlapped. In these CMDs the conspicuous MS is well populated from the oldest turnoff at $M_R=3.5$ up to the 0.03 Gyr isochrone. All the CMDs show a large fraction of young stars (< 1 Gyr old). The densely populated areas around the isochrones and the extension in luminosity of the Red Clump indicate a strong presence of intermediate-age stars. Lower panel shows the CMDs of the Western fields, located up to $\sim 2.9^\circ$ from the SMC center. In these fields, the intermediate-age population (~ 3 to 9 Gyr) is dominant. Fields *qj0036* and *qj0037* have a small fraction of stars born up to 0.1 Gyr ago, while *qj0033*, the most remote of the three, seems to have no stars younger than 1 Gyr and the main population is composed by stars between 6 and 9 Gyr old. Position angles (p), reddening ($E(B-R)$), metallicities and ages of the isochrones are labeled. A distance modulus $(m-M_0)=18.9$ has been assumed.

Fig. 2. Right panel showing the CMDs of the six Southern SMC fields. The same isochrones as in Figure 1 have been overlapped. In field *smc0057* a 0.03 Gyr isochrone with overshooting ($Z=0.004$) was also superimposed. This field is the closest to the SMC center and the one with the smallest position angle (in this Southern part), combination that gives a CMD with an important MS, where the 0.1 Gyr isochrone is still quite populated. The population gradient is depicted by the gradual lack of stars from 0.01 to 9 Gyr as going further South from the SMC center. Despite this, even in the more distant fields, the population is not purely old but dominated by intermediate-age stars as is shown in the zones around the 3, 4, 6, and 9 Gyr isochrones. Neither of the SMC CMDs in Figure 1 nor in this Figure present a blue horizontal branch. CMDs in both Figures are shown in order of increasing distance from the SMC center. Position angles (p), reddening ($E(B-R)$), metallicities and ages of the isochrones are labeled. A distance modulus $(m-M_0)=18.9$ has been assumed.