

UNIFORMITY DESPITE DIVERSITY: PN EXPANSION VELOCITIES IN THE LOCAL GROUP

M. G. Richer,¹ J. A. López,¹ M. G. Lee,² and N. Hwang²

We have undertaken a survey of planetary nebula (PN) expansion velocities in the [O III] λ 5007 line in Local Group galaxies visible from the OAN-SPM, Mexico using the Manchester Echelle Spectrograph (MES; Meaburn et al. 2003) on the 2.1-meter telescope. The instrumental profile has a velocity width of 11.5 km s^{-1} (2.7 pixel FWHM), and the nebular line profiles are always resolved.

The main results are: For the brightest of these spatially unresolved objects, asymmetries in the line profiles are clear. The mean expansion velocities (FWHM) seem remarkably uniform — despite the diversity of PN progenitors in different galaxies arising from very different stellar populations (Figure 1). Nevertheless, with larger samples, some differences are beginning to emerge. In M31, where we observe PNe sampling the bulge and different parts of the disk, a trend of decreasing expansion velocity with distance from the center is found. Statistically, the probability that the bulge and outer disk samples are drawn from populations with the same mean expansion velocity is only 3%. The PNe in M31’s outer and intermediate disk are drawn from the survey of Richer, Lee & Hwang (2004). NGC 205 and NGC 3109 stand out for having mean PN expansion velocities lower than the others. The O^{2+} zone could sample different parts of the nebular shell in objects arising from different stellar populations, but this is unlikely to be problematic since we observe the intrinsically brightest PNe in [O III] λ 5007, where the O^{2+} zone encompasses the majority of the mass.

MGR and JAL gratefully acknowledge the financial support of grants from CONACyT (CONACyT 43121) and UNAM (DGAPA IN 112103).




Fig. 1. For each galaxy, the point indicates the mean expansion velocity measured for its PN population; the error bars indicate the standard deviation of the expansion velocity distribution, and the numbers along the bottom axis indicate the number of PNe observed. We adopt the half-width at half-maximum as our measure of expansion velocity. Although this measure is arbitrary, most of the PNe have Gaussian line profiles at the S/N we achieve. The data for the Magellanic Clouds were taken from the literature (Dopita et al. 1985; Dopita et al. 1988), but analyzed in the same manner as the other data.

REFERENCES

- Dopita, M. A., Ford, H. C., Lawrence, C. J., & Webster, B. L. 1985, *ApJ*, 296, 390
 Dopita, M. A., Meatheringham, S. J., Webster, B. L., & Ford, H. C. 1988, *ApJ*, 327, 639
 Meaburn, J., López, J. A., Gutiérrez, L., Quirós, Murillo, J. M., Valdéz, J., & Pedrayes, M. 2003, *RevMexAA*, 39, 185
 Richer, M. G., Lee, M. G., & Hwang, N. 2004, *J. Korean Astron. Soc.*, 37, 269

¹Instituto de Astronomía, Universidad Nacional Autónoma de México, Apartado Postal 877, 22800 Ensenada, Baja California, México (richer@astrosen.unam.mx).

²Astronomy Program, SEES, BLD 25-1 Room 410, Seoul National University, San 56-1 Shillimdong, Gwanag-gu, Seoul 151-742, Korea.