

BLACK HOLE MASS ESTIMATES IN NEARBY AGN FROM HOST-BULGE PROPERTIES

E. Benítez,¹ A. Franco-Balderas,¹ I. Cruz-González,¹ L. López-Martin,² and V. H. Chavushyan³

In this work we present the results of an optical photometric study performed to seven nearby AGN that were selected with the SDSS-DR5 database. Surface photometric techniques have been applied to the data in order to obtain B/D ratio and therefore the $M_R(\text{Bulge})$ magnitudes. Bulge luminosities were then used to estimate the black-hole masse (M_{BH}) in these AGN.

The observations of nearby AGN has always been relevant since they enable us to study their host galaxies properties in a more detailed way. In recent years, it has been shown that host-bulge properties are correlated with the mass of the compact objects associated with the nuclei of normal and active galaxies (for a review on this topic, see Ferrarese & Ford 2005). Therefore, in order to understand the formation and evolution of black-holes (BH) it is necessary to estimate the M_{BH} of active galaxies over the widest possible ranges of host-galaxy types. This is the main goal of our study, still in progress.

In this work, we present the first results obtained from a study of 7 nearby AGN. We have selected objects flagged as QSO in the SDSS-DR5 that have a Petrosian-g magnitudes between $14 < M_g < 17$ and distances between $0.02 < z < 0.09$. An additional condition imposed to these AGN was that their host-galaxies should always have a major-axis size $> 15''$ in the isophote $\mu_B = 25 \text{ mag/arcsec}^{-2}$. These criteria allowed us to consider Type 1 (Sy1 and NLS1) as well as Type 2 (Sy2 and Sy1.9) AGN as good candidates for a deep imaging study using small-moderate size optical telescopes: NOT 2.5 m + ALFOSC, and SPM 1.5 m + Marconi. We present here the results obtained from our first couple of runs in (2006): (UT 1–2) with the NOT, and in November (UT 14–18) with the 1.5 m. We have observed Sy galaxies (i.e. with $M_B < -23$) that have a distance range between

TABLE 1
BLACK HOLE MASS

SDSS-Name	$M_R(\text{Bulge})$	$\log(M_{\text{BH}})$
J020615.98 – 001729.1	-21.96 ± 0.05	8.39 ± 0.49
J021011.49 – 090335.5	-20.03 ± 0.08	7.43 ± 0.48
J030417.77 + 002827.3	-20.83 ± 0.06	7.83 ± 0.48
J073106.86 + 392644.7	-19.67 ± 0.07	7.24 ± 0.48
J211646.34 + 110237.4	-20.50 ± 0.05	7.66 ± 0.48
J212851.19 – 010412.4	-18.78 ± 0.14	6.80 ± 0.48
J234428.81 + 134946.0	-19.03 ± 0.11	6.92 ± 0.48

174.89 and 340.65 Mpc ($H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.3$ and $\Omega_\lambda = 0.7$). We have obtained several broad-band B , V , R images with minimum exposure times of 2400 s in B , 1600 s in V and 800 s in R . Our images are deep enough since we want to have a $S/N \sim 3$ in the isophote $\mu_B = 25 \text{ mag/arcsec}^{-2}$. Although the nights assigned with the NOT telescope were not photometric, we got images with a seeing fluctuating from ~ 0.7 – $1.0''$. The average seeing measured at SPM in the R band was $\sim 1.2''$, but luckily the 4 assigned nights for this project were photometric dark nights. Thus, we have used the SPM data to calibrate the entire set of observed galaxies.

In order to perform the host-galaxy surface photometric study, we have followed the methodology given in Torrealba et al. (2006). Here we will present only part of our results. We have estimated the M_{BH} for each object using the correlation found by McLure & Dunlop (2002) but modified accordingly to our assumed cosmology. In Table 1 we show our estimates. These results, will be compared with M_{BH} estimations derived for the same AGN but with different methods.

REFERENCES

- Ferrarese, L., & Ford, H. 2005, Space Sci. Rev., 116, 523
 McLure, R. J., & Dunlop, J. S. 2002, MNRAS, 331, 795
 Torrealba, J., Benítez, E., Franco-Balderas, A., & Chavushyan, V. H. 2006, RevMexAA, 42, 3

¹Instituto de Astronomía, Universidad Nacional Autónoma de México, Apdo. Postal 70-264 México, D. F., México (erika@astroscu.unam.mx).

²Instituto de Astrofísica de Canarias, 38200, La Laguna, Tenerife, Spain (luislm@iac.es).

³Instituto Nacional de Astrofísica, Óptica y Electrónica, Apdo. Postal 51, 72000 Puebla, México (vahram@inaoep.mx).