

## STATISTICAL STUDY OF AXIAL ROTATION IN A TYPE STARS WITH NEW DETERMINATION OF $v \sin i$

M. Grosso<sup>1,2</sup> and H. Levato<sup>1,2</sup>

We have observed all A type stars of the southern hemisphere brighter than 6.5 mag in order to perform statistical studies of axial rotation. The spectra were obtained at the 2.15 m telescope at CASLEO. We have used the method of Diaz et al. (2011) to calculate axial rotation velocities.

Figure 1 shows the distribution of stars by spectral type including 66 Am and 42 Ap stars. Preliminary results for A6-A9 interval and the comparison with  $\langle v \sin i \rangle$  values from Royer et al. (2002, 2007) for a few common stars are shown in Figures 2 and 3. The last figure does not include Ap and Am stars.




Fig. 1. Distribution of Spectral Type for the programme stars.

### REFERENCES

- Diaz, C. G., Gonzalez, J. F., Levato, H., & Grosso, M. 2011, A&A, 531, A143  
 Royer, F., Grenier, S., Baylac, M. O., Gomez, A. E., & Zorec, J. 2002, A&A, 393, 897

<sup>1</sup>Instituto de Ciencias Astronómicas de la Tierra y del Espacio, CONICET, Casilla de Correo 49, CP 5400, San Juan, Argentina (mgrosso@icate-conicet.gob.ar).

<sup>2</sup>Universidad Nacional de San Juan, Argentina.




Fig. 2. Comparison with  $\langle v \sin i \rangle$  values from Royer.




Fig. 3. Distribution of  $\langle v \sin i \rangle$  values for this work.

Royer, F., Zorec, J., & Gomez, A. E. 2007, A&A, 463, 671