

H110 α AND C110 α EMISSION TOWARD THE PHOTODISSOCIATED REGION IN THE GGD14 COMPLEX

S. P. Treviño-Morales,¹ Y. Gómez,¹ C. A. Rodríguez-Rico,² G. Garay,³ L. F. Rodríguez,¹
and Á. Sánchez-Monge⁴

In this work we present high angular resolution observation of the C110 α radio recombination line (RRL) to study the photodissociated region (PDR) associated with the H II region of cometary morphology (VLA 1) in the GGD14 complex.

The GGD14 region is located in an active star-forming region embedded in the Monoceros molecular cloud at a distance of ~ 1 kpc (Rodríguez et al. 1980). This region contains an HII region (VLA 1) that seems to be ionized by a B0.5 ZAMS star (Gómez et al. 1998, 2000), a bipolar CO outflow (Rodríguez et al. 1982), and a cluster of water masers centered on the powering source of the molecular outflow (Tofani et al. 1995), which is located $\sim 30''$ to the northwest of the HII region. Gómez et al. 1998 has reported C92 α emission arising from an extended region of $\sim 20''$ in size that is closely associated with the HII region. The C92 α is interpreted as arising from a PDR around the HII region. Recently, Gómez et al. (2010) reported an expanding HI 21 cm champagne flow, with a size ten times larger than the HII region, suggesting that the neutral gas is expanding in a similar way as the ionized gas.

The H110 α and C110 α RRLs at 4.87 GHz were observed with the Very Large Array (VLA) in its C configuration. Data reduction was performed following the procedures for high frequency data of the software package AIPS. Both lines were detected, imaged and self-calibrated using the continuum emission, obtaining images with a synthesized beam of $7''.8 \times 5''.0$ at PA = -16° , and a rms noise of 1.41 mJy beam $^{-1}$ per 0.75 km s $^{-1}$ channel.

In Figure 1, we show the zero-order moment (integrated intensity) maps of both RRLs. We have spatially resolved the PDR surrounding the HII re-




Fig. 1. H110 α line emission (color scale) and C110 α line emission (white contours). Figure from Treviño-Morales, S., et al. (2011, in preparation).

gion in GGD14. We find that the C110 α line emission exhibits an extended arc structure that extends to the NW in between the HII region and the molecular gas, suggesting the presence of a high density PDR, while the HI 21 cm line emission extends from the NW to SE, likely a low density PDR (Gómez et al. 2010). This PDR structure is different to that expected from the Hollenbach & Tielens (1997) theoretical models.

In future work we will characterize the spectral line emission associated with the ionized gas, the PDR and the molecular cloud. The results obtained with this work could be a starting point for future research with ALMA.

REFERENCES

- Gómez, Y., et al. 2010, AJ, 140, 913
 Gómez, Y., et al. 1998, ApJ, 503, 297
 Gómez, Y., et al. 2000, RevMexAA, 36, 161
 Hollenbach, D. J., & Tielens, A. G. G. M. 1997, ARA&A, 35, 179
 Rodríguez, L. F., et al. 1982, ApJ, 260, 635
 Rodríguez, L. F., et al. 1980, ApJ, 235, 845
 Tofani, G., et al. 1995, A&AS, 112, 299

¹Centro de Radioastronomía y Astrofísica, Universidad Nacional Autónoma de México, Apdo. Postal 3-72, 58090 Morelia, Michoacán, Mexico (s.trevino@crya.unam.mx).

²Departamento de Astronomía, Universidad de Guanajuato, Guanajuato, Mexico.

³Departamento de Astronomía, Universidad de Chile, Santiago, Chile.

⁴Departament d'Astronomia i Meteorologia, Universitat de Barcelona, Spain.