RevMexAA (Serie de Conferencias), **40**, 44–45 (2011)

LONG-TERM OPTICAL PHOTOPOLARIMETRIC MONITORING OF BLAZARS AT SAN PEDRO MÁRTIR

E. Benítez,[1] J. Heidt,[2] D. Hiriart,[3] I. Agudo,[4,5] J. I. Cabrera,[6] D. Dultzin,[1] M. M. González,[1] J. M. López,[3] R. Mújica,[7] K. Nilsson,[8] R. Sacahui[1] and M. Sorcia[1]

RESUMEN

La variabilidad de luz polarizada en el óptico ha resultado ser una herramienta poderosa para estudiar a los Blázares debido a que permite determinar la intensidad y orientación del campo magnético asociado al jet relativista. También la luz polarizada es útil para estimar el sitio donde se origina la emisión de rayos-gama, aun cuando el mecanismo físico responsable de su producción sigue sin aclararse. Más importante aún, dado que las campañas en multifrecuencias típicamente se concentran en fuentes que están en estados de alta actividad, las características de las propiedades polarimétricas de Blázares en las fases pre- y post-estallido son menos conocidas. Por tanto, hemos iniciado un programa de monitoreo dedicado de Blázares de la muestra GASP en San Pedro Mártir para estudiar sus propiedades de variabilidad. En este trabajo queremos mostrar el estado en el que se encuentra el programa presentando algunos resultados preliminares sobre la variabilidad polarimétrica en el óptico y de rayos γ observada con Fermi-LAT en el blázar PKS 1510−089.

ABSTRACT

Variability of optical polarized light has result to be a powerful tool for studying Blazars, since it enable us to determine the strength and orientation of magnetic field associated with the relativistic jet. Also, polarized light is a valuable way of testing the location of the γ-ray emission, although the physical mechanisms responsible of its production remain unclear. Most importantly, since multi-wavelength campaigns typically concentrate on sources during highly active states, the characteristic polarimetric properties of Blazars in the pre-and post outbursts states are less known. Thus, we have started a dedicated monitoring program of Blazars from the GASP sample at San Pedro Mártir, in order to study their polarimetric variability properties. In this work we want to show the current status of the program presenting some preliminary results on the optical-polarimetric and γ-ray variability observed with Fermi-LAT on the blazar PKS 1510−089.

Key Words: galaxies: active — BL Lacertae objects: general

1. INTRODUCTION

Blazars are the extreme sub-population of the entire AGN class and include the BL Lac objects, and the flat spectrum radio quasars (FSRQ). Their overall spectral energy distribution (SED) shows two distinct emission peaks. Usually, the first peak is fitted assuming that the continuum emission is produced by synchrotron radiation, e.g., Marscher (1998). The second peak in the SED is usually observed in the high-frequency region (X-rays to γ-rays). The origin of the seed photons that are scattered in the second peak is still not completely well established. They may be synchrotron photons produced in the relativistic jet (SSC process) or can come from regions outside the jet (EC-models) like the accretion disk, the broad-line region or the dusty torus, e.g., Böttcher & Dermer (2010). Also, there are other models that assume that the γ-rays can be produced from particle cascades (Mannheim 1993; Blandford & Levinson 1995).

It has been recently shown that the study of the correlation between γ-ray activity and the variability of the optical polarization in blazars provides new information on the jet structure and the location and origin of the high-energy emission, see Agudo et al. (2011). Therefore, we are currently undertaken an optical-polarimetric monitoring program of 32 gamma-ray blazars (see Sorcia et al. 2011) that started in January 2008. In this paper, we will show our results on the object PKS 1510−089

[1]Instituto de Astronomía, Universidad Nacional Autónoma de México, Apdo. Postal 70-264, 04510 México D. F., Mexico (erika@astro.unam.mx).
[2]ZAH, Landesternwarte, Heidelberg, Germany.
[3]Instituto de Astronomía, UNAM, Ensenada BC, México.
[4]Institute for Astrophysical Research, Boston Univ., USA.
[5]Instituto de Astrofísica de Andalucía (CSIC), Spain.
[6]Instituto de Física, UNAM, México DF.
[7]Instituto Nacional de Astrofísica, Óptica y Electrónica, Puebla, México.
[8]Finnish Centre for Astronomy with ESO (FINCA), University of Turku, Finland.