RevMexAA (Serie de Conferencias), **40**, 181–182 (2011)

OBSERVATIONS OF THE PHOTODISSOCIATED HI REGION THAT SURROUNDS G213.880−11.837

C. A. Rodríguez-Rico,[1] Y. Gómez,[2] G. Garay,[3] C. Neria,[2] L. F. Rodríguez,[2] V. Escalante,[2] S. Lizano,[2] and M. Lebrón[4]

RESUMEN

Se presentan nuevas observaciones de la línea de HI a 21 cm hacia la región HII cometaria G213.880-11.837. Estas observaciones, llevadas a cabo a una resolución angular de ∼15″, revelan que el gas neutro en esta región es parte de un flujo en expansión. El análisis de la cinemática del gas HI sugiere que este gas está en una etapa de flujo de champagne: esto se basa en la diferencia que hay entre las velocidades radiales del gas neutro en la cola de la región (∼14 km s^{-1}) y la nube molecular ambiente (∼11.5 km s^{-1}) así como en la asimetría del gas de HI en emisión. Además de esta región fotodisociada, solamente hay otra región de este tipo que ha sido reportada presentando un flujo de champagne en HI (G111.61+0.37).

ABSTRACT

New observations of the HI 21 cm line toward the cometary HII region G213.880-11.837 are presented. These observations, carried out with an angular resolution of ∼15″, reveal that the neutral gas in this region is part of an expanding flow. The analysis of the kinematics of the HI gas suggests that the HI gas is undergoing a champagne flow: based on the difference between the radial velocities of the neutral gas in the tail (∼14 km s^{-1}) compared to the velocities of the ambient molecular gas (∼11.5 km s^{-1}) and the asymmetry of the HI emission. Besides this photodissociated region, there is only one other reported following an HI champagne flow (G111.61+0.37).

Key Words: HII regions — radio lines: ISM

1. GENERAL

GGD 14 is a star-forming region, part of the group GGD 12/13/14/15 embedded in the Monoceros molecular cloud at a distance of about 1 kpc (Rodríguez et al. 1980). GGD 14 contains a cluster of radio sources dominated by the compact (∼0.01 pc) cometary HII region (VLA 1; G213.880-11.837), excited by a B0.5 ZAMS star (Rodríguez et al. 1980; Kurtz et al. 1994; Gómez et al. 1998, 2000, 2002) and associated with IRAS 06084-0611. The ionized gas of the cometary HII region is undergoing a champagne flow with the head of the ionized champagne flow located to the NW and the extended emission toward the SE (Gómez et al. 1998). There is a velocity gradient along the major axis of the ionized champagne flow with increasing velocity from $V_{\rm LSR} \simeq 11$ km s^{-1} at the head of the flow to 15 km s^{-1} at the tail. As expected in a champagne flow the velocity at the head is very similar to the velocity of the ambient molecular gas ^{13}CO(2-1) which is ∼11.5 km s^{-1}(Qin et al. 2008). In addition of the ionized champagne flow, Gómez et al. (1998) reported the existence of an unresolved photodissociated region (PDR) surrounding the ionized gas.

2. RESULTS

The HI 21 line has been observed with the VLA (Gómez et al. 2010), at an angular resolution of 15″ and is detected both in absorption and emission in the velocity range from ∼1 to 20 km s^{-1} in agreement with previous results reported by Gómez et al. (1998). Figure 1 shows the integrated HI line emission and absorption image superposed on a near-IR continuum image at 4.5 μm from the Spitzer Space Telescope toward GGD 14. Our observations reveal that the size of HI line emission region is about a factor of ten larger than the HII region.

The HI optical depth was estimated using the HI absorption profile toward the HII region: $\tau_L(0) \simeq 5$. Using this result for the optical depth ($\tau_L(0) \gg 1$) the HI column density is estimated to be $N_{\rm HI} \geq 1.1 \times 10^{21}$ cm^{-2}. Further assuming that the physical

[1]Departamento de Astronomía, Universidad de Guanajuato, Callejón de Jalisco s/n, Col. Valenciana, Guanajuato, Guanajuato, Mexico (carlos@astro.ugto.mx).

[2]Centro de Radioastronomía y Astrofísica, Universidad Nacional Autónoma de México, Apdo. Postal 3-72, 58090 Morelia, Michoacán, Mexico (y.gomez@astrosmo.unam.mx).

[3]Departamento de Astronomía, Universidad de Chile, Casilla 36-D, Santiago, Chile (guido@das.uchile.cl).

[4]Departamento de Ciencias Físicas, Universidad de Puerto Rico, P.O. Box 23323, San Juan, PR 00931-3323, USA (mayra.lebron3@upr.edu).