RevMexAA (Serie de Conferencias), **40**, 227–228 (2011)

ATOMIC LINE WIDTHS BY THERMAL ENERGY FLUCTUATIONS IN STELLAR ATMOSPHERES

O. Cardona[1]

RESUMEN

Un método para encontrar los anchos de las líneas atómicas usando fluctuaciones de la energía térmica se presenta en este trabajo. Se hacen comparaciones con valores publicados experimentales y teóricos.

ABSTRACT

A method for finding the width of the atomic lines using the thermal energy fluctuations is presented in this work. Comparisons are made with published experimental and theoretical values.

Key Words: atomic processes — line: profiles — stars: atmospheres

1. INTRODUCTION

The broadening produced by the radiation of light by the atoms is called natural broadening and it occurs even for isolated atoms. The atoms embedded in a plasma will experience broadening of the lines caused by perturbations by other atoms or charged particles in the gas and is called pressure broadening. The impact and statistical quantum theories are local theories that explain the broadening (Griem 1997). The width at half maximum (FWHM) is used as a diagnostic tool in stellar atmospheres.

The method for finding the width of the atomic lines using the thermal energy fluctuations is presented in this work. Thermal energy fluctuations are a global collective phenomena by nature.

2. FLUCTUATIONS

The physical quantities which describe a macroscopic system in equilibrium are very nearly equal to their mean values. Nevertheless, fluctuations from the mean values occur. The probability of the fluctuations of the thermodynamic variables in a system is given by (Einstein 1907)

$$P(\Delta S) \; = \; Q \exp(\frac{\Delta S}{k}). \tag{1}$$

The mean square fluctuation of the energy σ^2, the variance, is given by (Kittel & Kroemer 1980)

$$\sigma^2 \; = \; \tau^2 (\frac{\partial U}{\partial \tau})_V, \tag{2}$$

where U is the mean energy of the thermodynamic system. The FWHM for the Gaussian distribution of the fluctuations is given by $\gamma = 2\sqrt{2 \ln 2}\sigma$. Equation (2) is the fundamental expression for developing the model.

[1]Instituto Nacional de Astrofísica, Óptica y Electrónica, Luis E. Erro 1, Tonantzintla, Puebla, 72840, Mexico (ocardona@inaoep.mx).

3. MODEL

The basic idea is that the change in energy of the levels is produced by the fluctuations in energy. The linear densities of the changes in energy and of the fluctuations are obtained. The principal assumption in this model is that these two linear energy densities are equal to each other. The linear density is used because the orbit is changed slightly by the perturbations, therefore, one has to take into account the size of the orbit to distribute those changes during the motion of the electron in the orbit.

3.1. *Hydrogenic Atoms*

Given the total number of particles per cubic centimeter N and the temperature T for a system the linear density of the fluctuations is obtained from the following expression (Cardona 2009)

$$\delta U = \frac{kT}{3L} = kT \frac{\sqrt[3]{N}}{3}, \tag{3}$$

and the linear density change in energy of the level is given by

$$\frac{\delta E_n}{2\pi r_n} = \frac{\alpha Ry}{\pi r_n n^3} \delta n, \tag{4}$$

where r_n is the radius of the orbit with principal quantum number n and α is a numerical factor to produces the FWHM of the level that now is a Gaussian distribution by the perturbations produced by the fluctuations. Our principal assumption requires equating equations (3) and (4) to obtain

$$\delta n = \frac{\pi r_n n^3}{\alpha} kT \sqrt[3]{N}. \tag{5}$$