RevMexAA (Serie de Conferencias), **40**, 239–240 (2011)

ON THE FORMATION OF THE MOST MASSIVE STARS IN THE GALAXY

R. Galván-Madrid,[1,2,3] Q. Zhang,[1] E. R. Keto,[1] L. F. Rodríguez,[2] and P. T. P. Ho[1,3]

RESUMEN

Presentamos resultados obtenidos con el Submillimeter Array (SMA) y el Very Large Array (VLA) en regiones de formación de estrellas masivas en sus primeras etapas de ionización. Las observaciones de gas ionizado y molecular con resolución angular de hasta $0.3''$ revelan rotación, colapso, y flujos eyectados en escalas de parsecs a < 0.05 pc. Los lugares con formación de estrellas tienen varias protoestrellas masivas en diferentes estados evolutivos. Los resultados indican que: (i) Las estrellas con $M_\star > 15\ M_\odot$ se forman por procesos de acreción similares a los de estrellas de baja masa, pero con ionización considerable. (ii) Los núcleos moleculares a partir de los cuales se forman estas estrellas continúan acretando gas de su ambiente. (iii) La acreción a la estrella puede continuar después del inicio de la ionización.

ABSTRACT

We present results of Submillimeter Array (SMA) and Very Large Array (VLA) studies of massive-star formation regions in the early stages of ionization. Molecular-line and ionized-gas observations at resolutions down to $0.3''$ reveal the presence of rotation, infall, and/or outflow from parsec scales to < 0.05 pc. The centers of star formation harbor small groups of massive (proto)stars at different evolutionary stages. These observations favor a picture in which: (i) Stars with $M_\star > 15\ M_\odot$ form by accretion processes similar to those of lower-mass stars, but with significant ionization. (ii) The molecular cores from which these stars form continue accreting from their environment. (iii) Accretion to the star can continue past the onset of ionization.

Key Words: H II regions — ISM: individual objects (W33A, G12.91–0.26, G20.08–0.14) — stars: formation

1. INTRODUCTION

To study the accretion dynamics over a range of scales in massive star formation regions (MSFRs), from the cluster scales (>1 pc) down to <0.05 pc, we set up a program of molecular line, recombination line, and continuum observations using the Submillimeter Array (SMA) and the Very Large Array (VLA) with several different angular resolutions. We chose exclusively sources with bolometric luminosities $L > 10^5\ L_\odot$ to ensure that they are forming (or can form) O-type stars ($M_\star > 15$ to 20 $M_\odot$). Here we summarize results of two sources representative of the low- and high-ionization ends of our sample. We refer the reader to Galván-Madrid et al. (2009, 2010) for further details.

2. W33A

W33A (G12.91–0.26, $L \approx 10^5\ L_\odot$) is a region with very high luminosity but very faint radio-continuum emission (Stier et al. 1984). We present the first resolved observations in both mm continuum and molecular-line emission of this MSFR. The two previously known mm cores (MM1 and MM2, see Figure 1) are fragmented into multiple peaks, and are at different evolutionary stages, as indicated by their molecular spectra, masses, temperatures, and continuum spectral indices. The warm-gas dynamics around the brightest core (MM1-Main at the center of MM1) indicates the presence of a circumstellar disk plus envelope rotating around a stellar mass of $M_\star \approx 10\ M_\odot$. MM1-Main also drives a powerful, high-velocity molecular outflow perpendicular to the disk. MM2, the coldest and most massive core, is not detected in hot-core lines but appears to drive a more modest outflow. Both MM1 and MM2 are located at the intersection of parsec-scale filamentary structures with a line-of-sight velocity offset of $\approx$2.6 km s^{-1}. The position-position-velocity structure of these filaments (see Figure 1) and a comparison with recent numerical simulations by Vázquez-Semadeni et al. (2009) suggest that star formation in W33A was triggered by the convergence of filaments of cold molecular gas.

3. G20.08N

G20.08-0.14 North (G20.08N, $L \approx 6.6 \times 10^5\ L_\odot$) was identified in the H II region VLA survey of Wood

[1]Harvard-Smithsonian Center for Astrophysics, USA (rgalvan@cfa.harvard.edu).

[2]Centro de Radioastronomía y Astrofísica, Universidad Nacional Autónoma de México, Mexico.

[3]Academia Sinica, Institute of Astronomy and Astrophysics, Taiwan.