

Fig. 1. K_s image of TR 14-N4 with the derived boundaries of the embedded cluster indicated by a circle. The centre of the field is at $\alpha = 10^{\rm h}\ 44^{\rm m}\ 31\overset{\rm s}{.}7$; $\delta = -59^\circ\ 33'\ 11''$ (J2000) and its size is $137'' \times 145''$. North is up, east is to the left.

ded in the dense cloud with the outflow directed close to the line of sight. We also update the near-infrared light curve (Figure 2) of the nearby embedded young objects #125 and #136, the latter known to have displayed K-band variations of more than four magnitudes (Tapia et al. 2003, 2006). It is evident that no clear variation pattern can yet be determined.



Fig. 2. Light curves in the K_s band, from 1993 November to 2010 April, of the young variable embedded stars (Tapia et al. 2006) Car I #136 (filled circles) and #125 (open circles). The photometric data of the comparison (non-variable) stars #91 (open triangles) and #142 (filled triangles) in the same field are also plotted.

The results describing the details of this work are being submitted to MNRAS.

MT acknowledges financial support from PAPIIT/UNAM grant IN 100210.

REFERENCES

Smith, N., & Brooks, J. K. 2008, Handbook of Southern Star Formation Regions Vol. 2, ed. B. Reipurth, (San Francisco: ASP), 138

Tapia, M., Roth, M., Vázquez, R. A., & Feinstein, A. 2003, MNRAS, 339, 44

Tapia, M., Persi, P., Bohigas, J., Roth, M., & Gómez, M. 2006, MNRAS, 367, 513