RevMexAA (Serie de Conferencias), **40**, 259–260 (2011)

INDUCED STAR FORMATION IN THE HII REGION SH2-54

J. Vasquez,[1,2] C. E. Cappa,[1,3] S. Cichowolski,[3,4] M. Ortega,[4] G. Romero,[3,5] and M. Rubio[6]

RESUMEN

En este trabajo presentamos un estudio multifrecuencia del medio interstelar en el entorno de la región HII Sh2-54. El gas molecular y el polvo se han analizado y descripto por medio de las observaciones milimétricas e infrarrojas, respectivamente. La emisión en 8.0 μm de *Spitzer*-IRAC presenta una compleja estructura filamentaria de gas fotodisociado, evidenciando la interacción entre la región HII y el gas molecular. Numerosos candidatos a objetos estelares jóvenes se encuentran proyectados sobre Sh2-54 y sus alrededores. Analizamos si la presencia de estos objetos puede ser explicada dentro de un escenario de formación estelar inducida.

ABSTRACT

We present a multiwavelength study of the interstellar medium around the HII region Sh2-54. The millimetric and infrared data allow us to analyze and characterize the molecular gas and dust components, respectively. The emision at 8.0 μm of *Spitzer*-IRAC shows a complex filamentary structure of photodissociated gas, revealing the interaction between the HII region and the molecular material. Several young stellar object (YSO) candidates have been detected projected towards Sh2-54 and its surroundings. We analyze if the presence of these YSO candidates can be explained by a triggered star formation scenario.

Key Words: HII regions — ISM: individual (Sh2-54) — stars: formation

1. GENERAL

The expansion of HII regions over molecular clouds generates potential sites of star formation. There are two main mechanisms that explain the induced star formation process on the edge of HII regions. The "collect and collapse" process (Elmegreen 1998) proposes the formation of massive stars, while the "radiatively driven implossion" model ("RDI", Lefloch & Lazareff 1994) focusses on the formation of low and intermediate mass stars, based on the compression of original overdensity regions. We can find in the literature a large number of star forming regions that can be explained by any of these two processes, e.g. *RCW 120* (Deharveng et al. 2009), *Gum 31* (Cappa et al. 2008), *SF 79* (Urquhart et al. 2004).

[1]IAR, CCT-La Plata, CONICET, C.C.5., 1894, Villa Elisa, Argentina.

[2]Cátedra de Espectroscopía Atómica y Molecular del Medio Interestelar, FCAG, UNLP, Argentina (jvasquez@fcaglp.unlp.edu.ar).

[3]Cátedra de Medio Interestelar, FCAG, UNLP, Argentina).

[4]IAFE (CONICET-UBA), Ciudad Universitaria, Buenos Aires, Argentina.

[5]Univ. de Valparaíso, Depto. de Física y Astronomía, Avenida Gran Bretaña 1111, Valparaíso, Chile.

[6]Depto. de Astronomía, Univ. de Chile, Casilla 36-D, Santiago, Chile.

2. SH2-54

Sh2-54 is an HII region of ∼6 arcmin in size, centered at (α, δ)(J2000)≃(18:17:50.0, –11:44:00.0), which is being ionized by several OB type stars. An DSS2 R image of the region is shown in the upper panel of Figure 1. It can be seen that the northwestern side of the nebula is surrounded by a high extinction region. The lower panel of Figure 1 shows the IRAC emission distribution at 8.0 μm with the integrated ^{12}CO(1-0) emission distribution (NANTEN data, HPBW = 2.7′ and 1 km s^{-1} velocity resolution), within the velocity range from +22 to +36 km s^{-1} (green contours) and the radio continuum emission distribution at 1465 MHz (VLA DnC conf. with a synthesized beam of 39.8×22.6 arcsec) indicated by blue contours. Both, the spectrophotometric and kinematical distances suggest for Sh2-54 a distance of $d = 2.0 \pm 0.5$ kpc. Based on the radio continuum and the millimetric emissions the main physical parameters of the ionized and molecular gas were estimated adopting an spherical geometry for the region (see Table 1).

3. STELLAR FORMATION

Based on the Allen et al.'s (2004) criteria, several YSO candidates projected onto the molecular emission were selected from the *Spitzer*-IRAC point source catalogue. The morphology of the phodissociation region (PDR) observed at 8 μm (see the lower